

FOR ANIMAL USE ONLY

**AIVLOSIN® WATER SOLUBLE POWDER**

Reg. No. G1598 Act 36/1947 N-SR0538

**CAUTION**

**STORAGE INSTRUCTIONS**

Store in a cool dry place.

**COMPOSITION**

Each 1 g of powder contains 850 mg of Acetyl-Isovaleryl-Tylosin tartrate (potency).

**INDICATIONS**

An aid in the prevention of Mycoplasmosis in broiler chickens and replacement pullets.

**WARNINGS**

- DO NOT SLAUGHTER BIRDS FOR HUMAN CONSUMPTION WITHIN 5 DAYS OF CESSATION OF TREATMENT.
- THIS DRUG MUST NOT BE USED IN LAYING HENS, WHERE THE EGGS ARE TO BE USED FOR HUMAN CONSUMPTION.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the distributor.

**PRECAUTIONS**

- This drug is a mild irritant and contact with the skin and eyes should be avoided.
- Wash thoroughly after handling the product.
- If accidental eye contact occurs rinse thoroughly with water.
- The medicated solution obtained after dilution of the product in water must be used immediately.
- Replace medicated water solutions daily.

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

• **Mixing**

**Prophylactic medication (broiler and layer pullets)**

Blend the contents of 1 sachet (25 g) into 2 l warm water.

Make up to 200 l of drinking water.

This is sufficient for treatment of 5 600 day-old broilers for 1 day or 2 800 chickens between 18 – 24 days old for 1 day, or 0,035 g/bird/day.

**Therapeutic medication (double strength)**

Blend the contents of 1 sachet (25 g) into 2 l warm water.

Make up to 100 l of drinking water.

• **Administration**

**Prophylactic Medication (Broiler and Layer Pullets)**

- Use only AIVLOSIN medicated water for 3 – 5 days post placement and again for 24 hours 3 - 4 weeks later.

- Replacement pullets may be treated for a further 24 – 48 hours (over and above the broiler programme) after management stress e.g. vaccination.

- This dosage rate is equivalent to approximately 75 g of AIVLOSIN per 500 kg or 1 000 live birds.

**Therapeutic Medication**

- Use only AIVLOSIN medicated water for at least 3 days when the birds show any signs of Chronic Respiratory Disease.

- This dosage rate is equivalent to ± 75 g of AIVLOSIN per 500 kg or 1 000 live birds.

**Registration holder:**

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 92/00835/07

P O Box 12406, Onderstepoort 0110

Tel: (012) 545-0004

SLEGS VIR DIERGEGBRUIK

**AIVLOSIN® WATER SOLUBLE POWDER**

Reg. No. G1598 Wet 36/1947 N-SR0538

**VERSIGTIG**

**BERGINGSANWYSINGS**

Bewaar in 'n koel droë plek.

**SAMESTELLING**

Elke 1 g poeier bevat 850 mg Acetiel-Isovaleriel-Tilosientartraat (sterkte).

**AANDUIDINGS**

'n Hulpmiddel vir die voorkoming van Kikoplasmose in braaikuikens en vervangingslêhennetjies.

**WAARSKUWINGS**

- MOENIE HOENDERS BINNE 5 DAE VANAF EINDE VAN DIE BEHANDELING VIR MENSlike VERBRUIK SLAG NIE.
- HIERDIE GENEESMIDDEL MOENIE IN LÊHENNE, WAAR EIERS VIR MENSlike VERBRUIK BEDOEL IS, GEBRUIK WORD NIE.
- Hou buite die bereik van kinders, oningeligte persone en diere.
  - Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

**VOORSORGMAATREËLS**

- Hierdie middel is matig irriterend en daarom moet kontak met die vel en oë vermy word.
- Was deeglik na hantering van die produk.
- As dit per ongeluk in aanraking kome met die oë, spoel deeglik met water.
- Die medisinale oplossing verkry na vermenging van die produk in water moet onmiddellik gebruik word.
- Vervang medisinale water daaglik.

**GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI**

• **Vermenging**

**Voorkomingsmedikasie (braaikuikens en vervangingslêhennetjies)**

Vermeng die inhoud van 1 pakkie (25 g) in 2 l warm water.

Maak op tot 200 l drinkwater.

Dit is genoeg vir die behandeling van 5 600 dagoud braaikuikens vir 1 dag of 2 800 hoenders tussen 18 – 24 dae oud vir 1 dag, of 0,035 g/kuiken/dag.

**Terapeutiese medikasie (dubbel sterkte)**

Vermeng die inhoud van 1 pakkie (25 g) in 2 l warm water.

Maak op tot 100 l drinkwater.

• **Toediening**

**Voorkomingsmedikasie**

- Gebruik AIVLOSIN medisinale water slegs vir 3 – 5 dae na plasing en weer vir 24 uur 3 - 4 weke later.

- Vervangingslêhennetjies kan behandel word vir 'n verdere 24 – 48 uur na enige bestuursspanning bv. inenting.

- Toediening soos hierbo aangedui is gelykstaande aan  $\pm$  75 g AIVLOSIN per 500 kg of 1 000 lewendige kuikens.

**Terapeutiese medikasie**

- Gebruik slegs AIVLOSIN medisinale water vir ten minste 3 dae wanneer die kuikens enige tekens van chroniese asemhalingsiekte (CAS) toon.
- Toediening soos hierbo aangedui is gelykstaande aan  $\pm$  75 g AIVLOSIN per 500 kg of 1 000 lewendige kuikens.

**Registrasiehouer:**

ECO Dieregesondheid Suider-Afrika (Edms) Bpk

Mpy. Reg. No. 92/00835/07

Posbus 12406, Onderstepoort 0110

Tel: (012) 545-0004