

FOR ANIMAL USE ONLY

AIVLOSIN® FG 50
Reg. No. G2951 Act36/1947
N-SR0918

An aid in the control of Mycoplasmosis in broiler chickens and replacement pullets. An aid in the control of swine enzootic pneumonia in pigs.

STORE IN A COOL DRY PLACE

COMPOSITION : 50 g acetyl-isovaleryl-tylosin tartrate per kilogram.

WARNINGS

- DO NOT SLAUGHTER BIRDS OR PIGS FOR HUMAN CONSUMPTION WITHIN 5 DAYS OF CESSATION OF TREATMENT.
- THIS DRUG MUST NOT BE USED IN LAYING HENS, WHERE THE EGGS ARE TO BE USED FOR HUMAN CONSUMPTION.
- Continuous administration of this drug for more than a week should be avoided.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Contact with the skin and eyes should be avoided.
- Wash thoroughly after handling the product.
- If accidental eye contact occurs, rinse thoroughly with water.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

• **Mixing**

• **Poultry**

AIVLOSIN FG 50 should be mixed into feed at a concentration of 200 ppm.
This is done by adding 4 kg per ton of AIVLOSIN FG 50.

• **Pigs**

AIVLOSIN FG 50 should be mixed into feed at a concentration of 20 – 50 ppm.
This is done by adding 0,4 kg to 1 kg per ton of AIVLOSIN FG 50.

The supplement must be completely dispersed throughout the total feed available to the animal at the recommended rate.
The required amount should first be mixed with ± 30 kg of feed, and then be thoroughly blended with the balance of the ration.

• **Treatment Guidelines**

For further information consult a veterinarian.

Presentation:25kg

Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd
Co. Reg. No. 92/00835/07
P O Box 12406, Onderstepoort 0110
Tel: (012) 545-0004