

APPROVED LABEL: CHLORTET 200 G – 25 KG (18.01.2003)

FOR ANIMAL USE ONLY

SLEGS VIR DIEREGERBUIK

CHLORTET 200 G

Reg. No. G3141 Act/Wet 36/100

For the treatment of respiratory and other infections caused by organisms susceptible to chlortetracycline in poultry, turkeys, pigs, calves and feedlot cattle.

Vir die behandeling van asemhalingskanaal- en ander infeksies veroorsaak deur organismes vatbaar vir chloortetrasiklien in pluimvee, kalkoene, varke, kalwers en voerkraalvee.

FEED ADDITIVE

STORAGE INSTRUCTIONS

Store in a cool, dry place.

COMPOSITION

Chlortetracycline 200 g/kg
(as the stable chlortetracycline calcium complex)

WARNINGS

- ALL FEED MEDICATED WITH CHLORTET 200 G MUST BE WITHDRAWN 7 DAYS BEFORE SLAUGHTER.
- DO NOT ADMINISTER TO LAYING HENS.
- DO NOT ADMINISTER TO BREEDING ANIMALS.
- NOT FOR DIRECT FEEDING. ACCURATE MIXING IS ESSENTIAL.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Wear protective clothing during addition of Chlortet 200 G to feed.
- Wash exposed skin after handling and before eating.
- Do not smoke or eat during use.

VOERBYVOEGSEL

BERGINGSAAWYSINGS

Bewaar in 'n koel, droë plek.

SAMESTELLING

Chloortetrasiklien 200 g/kg
(as die stabiele chloortetrasiklien kalsium kompleks)

WAARSKUWINGS

- ALLE VOER WAT MET CHLORTET 200 G BEHANDEL IS, MOET 7 DAE VOOR SLAGTYD ONTTREK WORD.
- MOENIE AAN LÊHOENDERS VOER NIE.
- MOENIE AAN TEELDIERE VOER NIE.
- MOENIE DIREK VOER NIE. AKKURATE VERMENGING IS NOODSAAKLIK.
- Hou buite die bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Dra beskermende klere gedurende toevoeging van Chlortet 200 G by voer.
- Was blootgestelde vel na hanteering en voor eet.
- Moenie rook of eet gedurende gebruik nie.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

The dosage rates given below are guidelines only – the inclusion rate should be determined by a veterinarian taking into account the incidence of disease and the severity of the challenge.

This supplement must be completely dispersed throughout the total feed available to the animal at the recommended rate.

To ensure uniform dispersion in the feed it is recommended that the correct amount of CHLORTET 200 G is first mixed with 10 X its weight of feed material before blending into the final feed.

GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

Die dosisse hieronder aangedui is slegs riglyne – toediening behoort deur 'n veearts vasgestel te word met inagneming van die siektetoestand en die erns daarvan.

Hierdie byvoegsel moet dwarsdeur die totale beskikbare voer teen die aanbevole vlak versprei word.

Om eweredige verspreiding te verseker word dit aanbeveel dat die aanbevole hoeveelheid CHLORTET 200 G eers met 10 X die hoeveelheid voer vermeng word voor dit in die finale rantsoen ingesluit word.

ANIMAL	Chlortetracycline activity g/ton Klortetrasiklien aktiewiteit g/ton	CHLORTET 200 G kg/ton	DIERSOORT
Poultry/turkeys	300 – 400	1,5 – 2,0	Pluimvee/kalkoene
Pigs	300 – 400	1,5 – 2,0	Varke
Feedlots: Cattle + calves	300 – 400	1,5 – 2,0	Voerkrale: Beeste + kalwers

These are average doses suitable for routine use. In the event of severe disease challenge, levels can be safely increased on the advice of a veterinarian to give up to 600 mg Chlortetracycline activity per kg of feed.

During treatment only Chlortetracycline-medicated feed should be used. Treatment should be continued for a period of 5 - 7 days or longer only if directed by a veterinarian.

Hierdie gemiddelde dosisse is toepaslik vir gewone gebruik. In geval van swaar siekte uitdaging kan die dosis met veiligheid voorhoog word met aanbeveling van 'n veearts tot 600 mg Chloortetrasiklien aktiewiteit per kg voer.

Gedurende behandeling moet slegs Chloortetrasiklien-behandelde voer gebruik word.

Behandel vir 5 – 7 dae of langer slegs op aanbeveling van 'n veearts.

CONTENTS 25 kg INHOUD

Lot No.:

Expiry/Verval:

Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd

ECO Dieregesondheid Suider-Afrika (Edms) Bpk

Co./Mpy. Reg. No. 92/00835/07

P O Box 12406, Onderstepoort, 0110

Tel: (012) 545-0004