

FOR EXTERNAL ANIMAL USE ONLY

ECOBASH CATTLE DIP

Reg. No. G3382 Act 36/1947

EMULSIFIABLE CONCENTRATE – CATTLE DIP AND SPRAY

- Controls ticks, has a detaching effect.
- Kills lice.
- Controls nuisance flies, biting flies, face flies,
- Protects against screw-worm infestation in cattle.

Contains: Cymiazol 17,5 % m/v; Cypermethrin 2,5 % m/v

STORAGE INSTRUCTIONS

- Store concentrate in original container away from food and feed.
- Store in a cool dry place.

WARNINGS

- NO MILK WITHDRAWAL.
- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST APPLICATION.
- Extremely toxic to fish.
- Handle with care. Poisonous by contact, swallowing or inhalation.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Avoid contact with concentrate. In case of accidental skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.
- Avoid prolonged contact with dipwash or inhalation or spray mist.
- Do not eat, drink or smoke while mixing or during use of dip.
- Prevent contamination of food, food utensils, grazing, drinking water, dams, rivers, etc.
- Destroy empty containers by perforation and flattening. Do not re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- DO NOT MIX WITH ANY OTHER DIP.
- SHAKE WELL BEFORE USE.
- Premix by adding the measured amount of concentrate slowly to double the volume of water and mix thoroughly by stirring.
- Add the premix to the bulk of water in the diptank or spray race sump. Make sure that in case of plunge tanks, the larger volume is poured in at the deep end.
- Mix contents of tank well before commencing dipping or spraying.

Dilution rates: Refer to table.

Recommendations:

- **Spray races:** Mix only enough spray-wash for one treatment and do not allow wash to stand over to the next spraying.
- **Boosting:** Boosting of spray wash is accomplished by adding 200 ml after every 100 head of cattle.
- **Hand spraying:** Thoroughly wet the animal to the skin, especially under the tail, the ears and between the legs where ticks cluster.

Dipping intervals:

- **Ticks, flies and screw-worm:** Dip or spray weekly.
- **Lice:** Dip or spray when necessary.

Dilution:

METHOD	FRESHFILL		REPLENISHMENT	
	ECOBASH CATTLE DIP	WATER	ECOBASH CATTLE DIP	WATER
Plunge dip	1,5 l	1 000 l	3 l	1 000 l
Spray race	1,5 l	1 000 l	1,5 l	1 000 l
Hand spraying	15 ml ECOBASH CATTLE DIP per 10 l water			

Presentation: 500 ml, 1 l, 5 l, and 20 l

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd
Co. Reg. No. 92/00835/07

Marketed by:

Afrivet Business Management (Pty) Ltd
Co Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004