

FOR ANIMAL USE ONLY

ECOLINT SUPER
Reg. No. G3065 Act 36/1947
N-SR0000

STORAGE INSTRUCTIONS

Store at room temperature below 25 °C.

INDICATIONS

A tapeworm remedy for cattle, sheep, goats and ostriches and a conical fluke remedy for cattle, sheep and goats.

COMPOSITION

Contains Resorantel 25 % m/v

WARNINGS

- DO NOT SLAUGHTER OR MILK FOR HUMAN CONSUMPTION FOR 2 DAYS AFTER TREATMENT.
- A transient fall in milk production in dairy cows may be expected after dosing.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED.

This stable suspension is ready for use without dilution or addition of any other substance.

- Sheep, goats and cattle: Dose orally 1 ml/4 kg body mass, for example:

| Body mass (kg) | Dose (ml) | No. of doses in pack | | |
|----------------|-----------|----------------------|--------|--------|
| | | 200 ml | 250 ml | 500 ml |
| 10 | 2,5 | 80 | 100 | 200 |
| 20 | 5,0 | 40 | 50 | 100 |
| 30 | 7,5 | 26 | 33 | 66 |
| 40 | 10,0 | 20 | 25 | 50 |
| 60 | 15,0 | 13 | 16 | 33 |
| 100 | 25,0 | 16 | 10 | 20 |
| 120 | 30,0 | 6 | 8 | 16 |
| 160 | 40,0 | 5 | 6 | 12 |
| 180 | 45,0 | 4 | 5 | 11 |

- Ostriches: Dose orally 1 ml/2 kg body mass, for example:

| Body mass (kg) | Dose (ml) | No. of doses in pack | | |
|----------------|-----------|----------------------|--------|--------|
| | | 200 ml | 250 ml | 500 ml |
| 5 | 2,5 | 80 | 100 | 200 |
| 10 | 5,0 | 40 | 50 | 100 |
| 30 | 15,0 | 13 | 16 | 33 |
| 50 | 25,0 | 8 | 10 | 20 |
| 70 | 35,0 | 5 | 7 | 14 |
| 100 | 50,0 | 4 | 5 | 10 |

EFFICACY

A. Tapeworms

| ANIMAL | TAPEWORM SPECIES | EFFICACY |
|-----------------|--|----------|
| Sheep and goats | <i>Moniezia expansa</i> (milk tapeworm) | Class I |
| | <i>Avitellina</i> spp. (narrow tapeworm) | Class I |
| | <i>Thysaniezia giardia</i> (frilled tapeworm) | Class I |
| Cattle | <i>Moniezia beneden</i> (milk tapeworm) | Class I |
| Ostriches | <i>Houttunyyia struthionis</i> (ostrich tapeworm) | Class I |

Explanation:

Class I: 100 % effective in more than 80 % of treated animals

Class II: 100 % effective in more than 60 % of treated animals.

Class III: 100 % effective in more than 50 % of treated animals.

Class X: Ineffective

B. Conical fluke

| ANIMAL | ADULT |
|-------------------------|---------|
| Sheep, goats and cattle | Class A |

Explanation:

Class A: 80 % effective in more than 80 % of treated animals.

Class B: 60 % effective in more than 60 % of treated animals.

Class C: 50 % effective in more than 50 % of treated animals.

Class X: Ineffective

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Presentation: 200 ml, 1lt, 5 lt.

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 1992/000835/07

P O Box 12406, Onderstepoort 0110

Tel: (012) 545-0004