

PROPOSED LABEL: OUTER CARTON - ECOMECTIN HORSE PASTE

FOR ANIMAL USE ONLY • SLEGS VIR DIEREGBRUIK

**ECOMECTIN**



**Horse Paste/Perdepasta**

Reg No. 3779 Act/Wet 36/1947

N-SR1458

**ANTIPARASITIC REMEDY FOR HORSES**

For use in the treatment and control of adult and immature roundworms, lungworms and bots of horses.

**ANTIPARASITIESE MIDDEL VIR PERDE**

Vir gebruik in die behandeling van volwasse- en onvolwasse rondewurms, longwurms en perdepapies.

**CAUTION/VERSIGTIG**

STORE BELOW 25 °C. PROTECT FROM LIGHT.

BERG ONDER 25 °C. BESKERM TEEN LIG.

**COMPOSITION/SAMESTELLING**

Ivermectin/Ivermektien 1,87 m/m %

Paste/Pasta 6,42 g

**FOR FULL PARTICULARS SEE ENCLOSED PACKAGE INSERT  
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET**

**1 Syringe/1 Spuit**

**Registration holder/Registrasiehouer:**

Eco Animal Health Southern Africa (Pty) Ltd  
Eco Dieregesondheid Suider-Afrika (Edms) Bpk  
Co./Mpy. Reg. No. 1992/000835/07  
P O Box 2009, Faerie Glen 0043, R.S.A.  
Tel: (012) 991 6416

† Macrocyclic lactone/Makrosikliese laktoon

Lot No:

Expires/Verval:

PROPOSED LABEL: SYRINGE  
ECOMECTIN HORSE PASTE

FOR ANIMAL USE ONLY • SLEGS VIR DIEREGERBRUIK

## **ECOMECTIN**

①†

### **Horse Paste/Perdepasta**

Reg No. 3779 Act/Wet 36/1947  
N-SR1458

#### **ANTIPARASITIC REMEDY FOR HORSES**

For use in the treatment and control of adult and immature roundworms, lungworms and bots of horses.

#### **ANTIPARASITIESE MIDDEL VIR PERDE**

Vir gebruik in die behandeling van volwasse- en onvolwasse rondewurms, longwurms en perdepapies.

### **CAUTION/VERSIGTIG**

STORE BELOW 25 °C. PROTECT FROM LIGHT.  
BERG ONDER 25 °C. BESKERM TEEN LIG.

#### **COMPOSITION/SAMESTELLING**

Ivermectin/Ivermektien 1,87 m/m %  
Paste/Pasta 6,42 g

**FOR FULL PARTICULARS SEE ENCLOSED PACKAGE INSERT  
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET**

**1 Syringe/1 Spuit**

#### **Registration holder/Registrasiehouer:**

Eco Animal Health Southern Africa (Pty) Ltd  
Eco Dieregesondheid Suider-Afrika (Edms) Bpk  
Co./Mpy. Reg. No. 1992/000835/07  
P O Box 2009, Faerie Glen 0043, R.S.A.  
Tel: (012) 991 6416

† Macrocyclic lactone/Makrosikliese laktoon

Lot No:  
Expires/Verval:

PROPOSED PACKAGE INSERT  
ECOMECTIN HORSE PASTE

FOR ANIMAL USE ONLY

**ECOMECTIC HORSE PASTE**



Reg. No. G3779 Act 36/1947  
N-SR1458

**Anitiparasitic remedy for horses**

For use in the treatment and control of adult and immature roundworms, lungworms and bots of horses.

**STORAGE INSTRUCTIONS**

Store below 25 °C.  
Protect from light.

**COMPOSITION**

Each syringe contains 6,42 g paste which contains 1,87 m/m % Ivermectin.

**WARNINGS**

- WITHDRAWAL PERIOD 21 DAYS.
- Ecomectin Horse Paste has been formulated specifically for use in horses only.
- Dogs and cats may be adversely affected by the concentration of the ivermectin in this product if they are allowed to ingest spilled paste or have access to the used syringes.
- Some horses carrying heavy infection of *Onchocerca* microfilariae have experienced oedema and pruritus following dosing, assumed to be the result of death of large numbers of microfilariae. These signs resolve within a few days but symptomatic treatment may be advisable.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

**PRECAUTIONS**

- EXTREMELY DANGEROUS TO FISH AND AQUATIC LIFE. Do not contaminate surface waters or ditches with product or used container.
- Drug containers and any residual contents should be disposed of in a safe manner. Syringes must not be re-used for any other purpose.
- Do not smoke, eat or drink while handling the product.
- Wash hands after using the product

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

**Dosage**

One syringe division of paste per 100 kg body mass (based on a recommended dosage of 200 µg ivermectin per kg body mass).

The total contents of one syringe (six divisions) will dose one 600 kg horse.

## Administration

The paste is given by the oral route.

The syringe should be inserted at the interdental space and the paste deposited on the base of the tongue.

Ecomectin Horse Paste can be administered to mares at any stage during pregnancy or lactation.

Ecomectin Horse Paste can be used in breeding mares and stallions and will not affect fertility.

Ecomectin Horse Paste can be given to all ages of animals including young foals.

## EFFICACY

Ecomectin Horse Paste at the recommended dose rate of 200 µg ivermectin per kg body mass provides effective treatment and control of the following parasites of horses:

Worm species	Immature	Adult
<b>Large strongyles</b>		
<i>Strongylus vulgaris</i> (and arterial phase)		*
<i>S. edentatus</i> (and tissue phase)		*
<i>S. equinus</i>		*
<i>Tridontophorus</i> spp.		*
<b>Small strongyles</b> (including benzimidazole resistant strains)		*
<i>Oxyuris equi</i>	*	*
<i>Parascaris equorum</i>		*
<i>Trichostrongylus axei</i>		*
<i>Habronema muscae</i>		*
<i>Strongyloides westeri</i>		*
* Microfilariae of <i>Onchocerca</i> spp.		
* Oral and gastric stages of bots, <i>Gastrophilus</i> spp.		

\* Control =  $\geq 90$  % effective.

## Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd

ECO Dieregesondheid Suider Afrika (Edms) Bpk

Co./Mpy. Reg. No. 1992/000835/07

P O Box 2009, Faerie Glen 0043, R.S.A.

Tel: (012) 991 6416

† Macrocyclic lactone

VOORGESTELDE VOUBILJET  
ECOMECTIN HORSE PASTE

SLEGS VIR DIEREGERBRUIK

**ECOMECTIN PERDEPASTA**

Reg. No. G3779 Wet 36/1947  
N-SR1458



**Antiparasitiese middel vir perde**

Vir gebruik in die behandeling van volwasse- en onvolwasse rondewurms, longwurms en perdepapies.

**BERGINGSAAWYSINGS**

Berg onder 25 °C.  
Beskerm teen lig.

**SAAMESTELLING**

Elke spuit bevat 6,42 g pasta met 1,87 m/m % Ivermektien.

**WAARSKUWINGS**

- ONTTREKINGSPERIODE 21 DAE.
- Ecomectin Perdepasta is slegs vir gebruik in perde geformuleer.
- Nuwe effekte mag voorkom in honde en katte as hulle in aanraking kom met die Ivermektien in die produk, hetsy deur inname van afval of gebruikte spuite.
- In perde met 'n swaar infestasië van *Onchocera* mikrofilaria kan edeem en jeuk voorkom na doseering, waarskynlik as 'n gevolg van groot getalle mikrofilariae sterftes. Die simptome sal gewoonlik oor 'n paar dae verdwyn maar simptomatiese behandeling mag aanbeveel word.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

**VOORSORGMAATREËLS**

- UITERS GIFTIG VIR VIS EN WATERLEWE. Moenie water oppervlaktes of slote met produk of gebruikte houers besoedel nie.
- Wegdoening van houers en enige oorblywende inhoud moet op 'n veilige manier geskied word.
- Moenie rook, eet of drink terwyl produk gehanteer word nie.
- Was hande na hantering van produk.

**GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI**

**Dosis**

Een spuitafdeling van pasta per 100 kg liggaamsmassa (gebaseer op 'n aanbevole dosis van 200 µg ivermektien per kg liggaamsmassa).

Die totale inhoud van een spuit (ses afdelings) sal een 600 kg perd doseer.

## Toediening

Die pasta word oraal gedoseer.

Die spuit moet by die interdendale opening toegedien word en die pasta aan die basis van die tong gedeponeer word.

Ecomectin Perdepasta kan tydens enige stadium van dragtigheid of laktasie aan merries toegedien word.

Ecomectin Perdepasta kan in teelmerries en hingse gebruik word sonder om vrugbaarheid te affekteer.

Ecomectin Perdepasta kan toegedien word aan diere van enige ouderdom, insluitend jong vulle.

## DOELTREFFENDHEID

Ecomectin Perdepasta teen die aanbevole dosis van 200 µg ivermektien per kg liggaamsmassa bewerkstellig effektiewe behandeling en beheer van die volgende parasiete van perde:

Wurmspesie	Onvolwasse	Volwasse
<b>Groot strongiele</b> <i>Strongylus vulgaris</i> (en bloedvat fase) <i>S. edentatus</i> (en weefsel stadium) <i>S. equinus</i> <i>Tridontophorus</i> spp.		* * * *
<b>Klein strongiele</b> (insluitend bensimidazool weerstandbiedende stamme)		*
<i>Oxyuris equi</i> <i>Parascaris equorum</i> <i>Trichostrongylus axei</i> <i>Habronema muscae</i> <i>Strongyloides westeri</i>	*	* * * * *
* Mikrofilaria van <i>Onchocerca</i> spp. * Orale- en buikstadia van perdepapie, <i>Gastrophilus</i> spp.		

\* Beheer =  $\geq 90$  % effektief

## Registrasiehouer:

ECO Dieregesondheid Suider-Afrika (Edms) Bpk

Mpy. Reg. No. 1992/000835/07

P O Box 2009, Faerie Glen 0043, R.S.A.

Tel: (012) 991 6416

† Makrosikliese laktoon