

FOR ANIMAL USE ONLY

ECOMINTIC 50

Reg. No. G2000 Act 36/1947

STORAGE INSTRUCTIONS

Store in a cool, dry place.

COMPOSITION

Fenbendazole 5 % m/v.

INDICATIONS

A remedy against roundworms in cattle, horses and ostriches and roundworms and milk tapeworm in sheep and goats.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF TREATMENT.
- Do not mix or dilute with any other substance.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large range of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- It is recommended that sheep be vaccinated against pulpy kidney disease before any worm remedy is administered.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

Use as an oral drench.

Shake well before use.

Dosage rate:

Cattle: 1 ml/10 kg body mass.

Horses: 2 ml/10 kg body mass.

Sheep and Goats:

• **Roundworms** 1 ml/10 kg body mass.

• **Milk tapeworm (*M. expansa*)** 2 ml/10 kg body mass.

Ostriches: 3 ml/10 kg body mass.

Example for cattle and horses (roundworms):

Body mass (kg)	Dose (ml)		No. of doses in pack			
	Cattle	Horses	Cattle		Horses	
			200 ml	500 ml	200 ml	500 ml
50	5	10	40	100	20	50
100	10	20	20	50	10	25
150	15	30	13	33	6	16
200	20	40	10	25	5	12
250	25	50	8	20	4	10
300	30	60	6	16	3	8
350	35	70	5	14	2	7

Example for sheep, goats and ostriches (roundworms and milk tapeworm):

Body mass (kg)	Dose (ml)			No. of doses in pack					
	Sheep and Goats		Ostriches	Sheep and Goats				Ostriches	
	Roundworm	Milk Tapeworm (<i>M. expansa</i>)		Roundworm		Milk Tapeworm (<i>M. expansa</i>)			
				200 ml	500 ml	200 ml	500 ml	200 ml	500 ml
10	1	2	3	200	500	100	250	66	166
20	2	4	6	100	250	50	125	33	83
30	3	6	9	66	166	33	83	22	55
40	4	8	12	50	125	25	62	16	41
50	5	10	15	40	100	20	50	13	33

EFFICACY

- **Cattle, sheep and goats**

Worm species	Immature	Adult
• Cattle		
Wireworm	*	*
Brown stomachworm	*	*
Nodular worm	*	*

Hookworm	*	*
Lungworm	**	*
• Sheep and Goats		
Wireworm	*	*
Brown stomachworm	*	*
Bankruptworm	*	*
Long-necked bankruptworm	*	*
White bankruptworm	*	*
Hookworm	*	*
Large-mouthed bowelworm	*	*
Nodular worm	*	*
Lungworm	*	*
Milk tapeworm (<i>Moniezia expansa</i>)	Class I	

* Control = ≥ 90 % effective

** Aids in Control = 60 – 89 % effective

Class I: 100 % effective in more than 80 % of treated animals.

Kills worm eggs present in the animal at the time of dosing.

• Horses

* Bloodworms (large strongyles), redworms (small strongyles), pinworm (*Oxyuris*), ascarids and stomach worm (*Habronema majus*).

• Ostriches

* *Libyostrongylus douglassii*

* Control = ≥ 90 % effective

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

† Benzimidazole

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 92/00835/07

P O Box 12406, Onderstepoort, 0110

Tel: (012) 545-0004