

FOR ANIMAL USE ONLY

ECOX® 200 MG
Reg No. G3219 Act 36/1947

PREMIX

An aid in the prevention of coccidiosis in broiler chickens, chickens reared for laying and turkeys up to 16 weeks of age. For improved feed efficiency in feedlot cattle.

STORAGE: Store in a cool dry place. Close container after use. Store away from feed and foodstuffs.

Contains: Each kilogram of premix contains monensin sodium equivalent to 200 g monensin activity.

WARNINGS

- DO NOT FEED TO HORSES OR OTHER EQUINES, IT MAY PROVE FATAL.
- DO NOT FEED TO LAYING POULTRY, DAIRY CATTLE AND MILK GOATS.
- Animals should not be treated with products containing tiamulin or troleandomycin while receiving, or for at least 7 days before or after receiving, feed containing ECOX 200 MG. Severe growth depression or death may result.
- This product should not be mixed with another coccidiostat.
- Do not exceed the recommended levels of ECOX 200 MG premix as reduced average daily gain may result.
- Handle with care. Poisonous when swallowed.
- Avoid skin contact and inhalation.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Operators handling ECOX 200 MG should avoid direct contact with the skin and should wear protective clothing, gloves and a dust mask when preparing medicated feed.
- Wash thoroughly after handling the product.
- If accidental eye contact occurs, rinse thoroughly with water.
- **First aid treatment:** In case of poisoning or ingestion, call a doctor and make this information available. If swallowed, induce vomiting by tickling the back of the throat and repeat until vomit is clear.
- **Note to doctor:** Perform gastric lavage if the patient has not vomited and treat symptomatically.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

It is recommended that ECOX 200 MG should be thoroughly mixed in a small quantity of feed (20 – 30 kg) before it is incorporated into the bulk of the total feed to be prepared.

Broilers	ECOX 200 MG should be mixed into feed at a rate of 500 – 600 g per metric ton, to provide a level of 100 – 120 mg/kg Monensin activity in finished feed. This feed should be fed continuously as the only ration.
Chickens reared for laying/	ECOX 200 MG should be mixed into feed at a rate of 500 – 600 g per metric ton, to provide a level of 100 – 200 mg/kg Monensin activity in finished feed. Feed continuously until 16 weeks of age.
Turkeys	ECOX 200 MG should be mixed into feed at a rate of 450 – 500 g per metric ton, to provide a level of 90 – 100 mg/kg Monensin activity in finished feed. Feed continuously until 16 weeks of age.
Feedlot cattle	ECOX 200 MG should be mixed into feed at a rate of 50 – 150 g per metric tone, to provide not less than 10 g nor more than 30 g Monensin per metric ton of total ration, so that each animal receives not less than 50 mg nor more than 300 g Monensin per head per day.
	Rations containing silage or other wet feeds should be corrected to a 90 % dry matter basis.

Complete cattle feedlot feeds

Active ingredient per metric ton (ppm)/	*Consumption Monensin/head/day (mg)/	Gram ECOX 200 MG premix per metric ton/
10	90	50
15	135	75
20	180	100
30	270	150

* At 9 kg intake per head of final ration.

Presentation: 20 kg (25 kg)

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 1992/000835/07

P O Box 12406, Onderstepoort 0110

Tel: (012) 545-0004