

FOR ANIMAL USE ONLY

ERADIWORM

Reg. No. G0000 Act/Wet 36/1947

Roundworm and tapeworm remedy for sheep, goats, cattle and ostriches.

STORE IN A COOL, DRY PLACE.

Contains: Levamisole HCl 37,5 mg/ml ; Praziquantel 18,8 mg/ml

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF TREATMENT.
- No worm remedy should be dosed to sheep and goats that have not been immunised against pulpy kidney.
- When treated with levamisole, the odd animal may develop an allergic-type reaction of a transient nature which normally disappears within 24 hours. An antihistamine may be administered. Under no circumstances must cortisone preparations be used.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- **Sheep and goats:** 2 ml/10 kg body mass.
- **Cattle:** 10 ml/50 kg body mass.
- **Ostriches:** 1 ml/1 kg body mass.

EFFICACY

Worm species	Sheep/goats		Cattle	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Brown stomachworm		**		**
Long-neck bankruptworm	**	*		
Large-mouthed bowelworm	*	*		
Nodular worm	*	*	**	*
Lungworm	**	**		
Hookworm	*	*		*
Bankruptworm (<i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm (<i>Cooperia</i> spp.)			*	*
Milk tapeworm	CLASS 1			
			Ostriches	
Wireworm (<i>Libostrongylus douglassi</i>)			**	*
Hookworm (<i>Codiostomum struthionis</i>)			**	*
Tapeworm (<i>Houttuynia struthionis</i>)			**	*

* Controls = \geq 90 % effective
** Aids in control = 60 – 89 % effective
Class1: 100 % effective in more than 80 % of treated animals.

200 ml (250 ml, 500 ml, 1 l, 5 l, 10 l)

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 92/00835/07

P O Box 12406, Onderstepoort, 0110

Tel: + 27 (0)12 545-0004