

FOR ANIMAL USE ONLY

SULPHAMEZATHINE® 16 %
Reg. No. G0802 (Act 36/1947)
N-SR0855

INDICATIONS

For the treatment of coccidiosis and coryza (croup) in poultry, and coccidiosis in calves, lambs, kids and rabbits.

STORAGE INSTRUCTIONS: Store in a cool dry place.

COMPOSITION: Sulphadimidine sodium BP 16 % m/v.

WARNINGS

- ALLOW 7 DAYS BETWEEN LAST TREATMENT AND SLAUGHTER FOR HUMAN CONSUMPTION.
- EGGS LAID DURING TREATMENT AND FOR THREE DAYS THEREAFTER, MUST NOT BE USED FOR HUMAN CONSUMPTION.
- THE USE OF THIS REMEDY, OR ANY OTHER SULPHONAMIDE, MAY CAUSE A TEMPORARY DROP IN EGG PRODUCTION IN LAYING HENS.
- **N.B.** Animals must have free access to ample drinking water during and for at least 2 days after last treatment.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
FRESH SOLUTIONS MUST BE PREPARED DAILY.

Dosage

• **Rabbits and poultry**

13 ml/1 l of drinking water.

Administer for 3 days, withhold for 2 days then administer for a further 3 days.

No other drinking water or succulent green feed is allowed during the days of treatment.

• **Lambs, kids and calves**

Initial dose of 14 ml/10 kg body mass orally.

Maintenance dose of 7 ml/10 kg for a further 2 days.

If further treatment seems necessary consult your veterinarian.

Presentation: 100 ml and 500ml

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004