

FOR ANIMAL USE ONLY

TRAMISOL® PLUS
Reg. No. G1830 Act/Wet 36/1947
N-SR0883

Roundworm- and liverfluke remedy for cattle, sheep and goats as well as a nasal worm remedy for sheep and goats.

Contains: Levamisole hydrochloride 2,5 % m/v; Rafoxanide 2,5 % m/v

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 28 DAYS OF DOSING.
- DO NOT DOSE LACTATING ANIMALS WHOSE MILK OR MILK PRODUCTS ARE INTENDED FOR HUMAN CONSUMPTION.
- MILK FROM TREATED ANIMALS MUST NOT BE USED FOR HUMAN CONSUMPTION.
- Care must be taken in calculating the correct dosage rate as overdoses may lead to toxicity.
- Always ensure that all drenching procedures are carefully carried out. Care should be exercised in the dosing and handling of animals.
- No worm remedy, not even Tramisol Plus, should be dosed to sheep and goats that have not been immunised against Pulpy Kidney (enterotoxaemia).
- In case of poisoning call a doctor and make this information available.
- Keep out of reach of children, uninformed persons and animals.
- Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

SHAKE WELL BEFORE USE.

Do not treat animals more frequently than every 3 weeks.

• **Dosage**

Sheep and goats: 3 ml per 10 kg body mass.

Cattle: 15 ml per 50 kg body mass.

Example: Cattle, sheep and goats

Body mass (kg)	Dose (ml)	No. of doses in pack			
		200 ml	1 l	5 l	10 l
10	3	66	333	1 666	3 333
20	6	33	166	833	1 666
30	9	22	111	555	1 111
40	12	16	83	416	833
50	15	13	66	333	666
100	30	6	33	166	333

• **Efficacy**

ROUNDWORMS	Sheep and goats		Cattle	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Brown stomachworm	*	*		
Long-necked bankruptworm	*	*		
Large-mouthed bowelworm	*	*		
Nodular worm	*	*	**	*
Lungworm	*	*	*	*
Hookworm	*	*		*
Bankruptworm (<i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm (<i>Cooperia</i> spp.)			*	*
FLUKES				
Liver fluke (<i>Fasciola hepatica</i>)	6 weeks **	*		*
Giant liver fluke (<i>Fasciola gigantica</i>)	8 weeks *	*		*
NASAL WORM				
<i>Oestrus ovis</i>	* (all larval stages)			

* Controls = ≥ 90 % effective

** Aids in control = 60 – 89 % effective

The list contains the more important endoparasitic species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

MAIN PANEL LABEL: 200 ml, (1 l, 5 l, 10 l)

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545 0004