

FOR ANIMAL USE ONLY

TRAMIZAN®
Reg. No. G0809 Act 36/1947
N-SR0884

Roundworm, liver- and conical fluke remedy for cattle, sheep and goats.

Contains: Levamisole hydrochloride 2,5 % m/v; Oxytocosanide 3,4 % m/v

WARNINGS

- ALLOW 7 DAYS BETWEEN LAST TREATMENT AND SLAUGHTER FOR HUMAN CONSUMPTION.
- No worm remedy, not even Tramizan, should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxaemia).
- In a low percentage of high-producing dairy cows the following transient reactions may be noted: diarrhoea, salivation and swelling of the head.
- In case of poisoning call a doctor and make this information available.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

SHAKE WELL BEFORE USE.

Tramizan may be dosed to all cattle, sheep and goats without any pre-dosing starving or change of diet.

• **Dosage**

Cattle: 15 ml/50 kg body mass.

Body mass (kg)	Dose (ml)	No. of doses in pack		
		1 l	5 l	10 l
50	15	66	333	666
150	45	22	111	222
250	75	13	66	133
300	90	11	55	111
400	120	8	41	83
500	150	6	33	66

Sheep and goats: 3 ml/10 kg body mass.

Body mass (kg)	Dose (ml)	No. of doses in pack		
		1 l	5 l	10 l
0 – 10	3	333	1 666	3 333
11 – 20	6	166	833	1 666
21 – 30	9	111	555	1 111
31 – 40	12	83	416	833
41 – 50	15	66	333	666
51 – 60	18	33	277	555

Cattle, sheep and goats (*Paramphistomum* spp.):

3 ml/10 kg body mass followed by a second dose 72 hours later.

• **Efficacy**

ROUNDWORMS	Sheep and goats		Cattle	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Brown stomachworm	**	**		
Long-necked bankruptworm	*	*		
Large-mouthed bowelworm	*	*		
Nodular worm	*	*	**	*
Lungworm	**	*	**	
Hookworm	*	*		*
Bankruptworm (<i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm (<i>Cooperia</i> spp.)			*	*
	Sheep and goats		Cattle	
	Immature	Adult	Immature	Adult
FLUKES				
Liver fluke (<i>Fasciola hepatica</i>)		*		*
Giant liver fluke (<i>Fasciola gigantica</i>)		*		*
Conical fluke (<i>Paramphistomum</i> spp.)	*	*	*	*

* Controls = ≥ 90 % effective

** Aids in control = 60 – 89 % effective

The list contains the more important endoparasitic species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Presentation: 200 ml (1 l, 5 l, 10 l)

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004