

FOR EXTERNAL ANIMAL USE ONLY

TRIATIX 125
Emulsifiable concentrate

Reg. No. G3189 Act 36/1947 N-SR1081

OXPECKER COMPATIBLE

STORAGE INSTRUCTIONS

Protect product from extremes of heat and cold and from dampness.

COMPOSITION

Amitraz 12,5 % m/v

INDICATIONS

CATTLE

Controls ticks. Kills lice and mange mites.

SHEEP AND GOATS

Controls ticks. Controls sheep scab. Kills itch mites and goat mange mites.

WARNINGS

• **WITHDRAWAL PERIOD:**

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST DIPPING.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this label available to him.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Tightly reseal the container if all the contents have not been used.
- Destroy empty containers.
- Wash hands thoroughly after use.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

For external use on cattle, sheep and goats.

TRIATIX 125 should not be mixed with any other dip.

CATTLE

• **Spraying**

Dipwash should be freshly prepared for each spraying.

Add required amount of TRIATIX 125 to ± 10 l water in a suitable container.

When well mixed add to measured bulk water in spray sump and stir thoroughly.

Use at a rate as per the table below:

TRIATIX 125	WATER
100 ml	50 l
200 ml	100 l
250 ml	125 l
500 ml	250 l
1 l	500 l
5 l	2 500 l
10 l	5 000 l
20 l	10 000 l

Boost

It is recommended that not more than 400 head be sprayed in the spray race at one time, especially where fouling is a problem. If, however, this figure is exceeded, boosting at the rate of 100 ml per 100 head, commencing at 400 head, can be undertaken to maintain a satisfactory deposit of active ingredient, especially where tick-borne diseases are a problem.

• **Plunge dip**

Total replenishment: This requires that the diptank is topped up to the desired level on each dipping occasion. TRIATIX 125 is then added on every occasion at a rate of 1 l to 2 500 l water, taking the total amount of water in the tank into the calculation.

Lime stabilised:

- **Fresh fill** Add TRIATIX 125 at the rate of 1 l per 500 l water.

At the end of each dipping add 5 kg calcium hydroxide (75 %) per 100 head of cattle dipped.

- **Replenishment** Add TRIATIX 125 at the rate of 1 l per 500 l water added.

At the end of each dipping, calcium hydroxide (75 %) must be added to the tank at a rate of 5 kg per 100 head of cattle dipped to ensure that the dipwash pH is in the region of 12.

Frequency:

- **Ticks and mange** Spray/dip weekly
- **Lice** Spray/dip twice with a 7 day interval

Should dipping intervals extend beyond 3 weeks, additional lime should be added at a rate of 1 kg lime (75 % active) per 1 000 l water in the dip tank. The dipwash should be adequately protected for 2 months if this procedure is followed.

SHEEP AND GOATS

TRIATIX 125 should be administered by means of either:

1. 2 Dippings using fresh, unstabilised dipwash. TRIATIX 125 should be added at a rate of 1 l per 375 l water, taking the total amount of water in the tank into consideration.
OR
2. A fresh, unstabilised dipwash for the first dipping, which is then stabilised with the addition of lime at a rate of 4 kg lime per 750 l dipwash, added immediately after the initial dipping. This stabilised dipwash is then suitable for the second dipping, 7 days later.

Frequency:

Sheep and goats should be dipped twice with a 7 day interval to kill sheep scab, itch and goat mange mites. Ensure that the animals are immersed for approximately 1 minute with the heads being submerged at least 3 times in order to ensure that the inside of the ears are thoroughly wetted.

N.B. Sheep scab is a notifiable disease and must be reported to the nearest State Veterinarian.

For the control of bacterial build-up and for the prevention of post-dipping lameness:

COPPER SULPHATE can be added at a rate of 500 g per 1 000 l of water.

OR

ZINC SULPHATE can be added at a rate of 1 kg/1 000 l of dipwash.

For the control of post-dipping lameness and lumpy wool:

ZINC SULPHATE can be added at a rate of 1 kg/400 l of dipwash.

For full details see labels of the abovementioned products.

Presentation: 200 ml, 500 ml, 1 l, 5 l, 20 l

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004