

APPROVED LABEL: TYLECO FG 100 (G2152)

FOR ANIMAL USE ONLY

**TYLECO F.G. 100**

Reg. No. G2152 Act 36/1947

An aid in improving growth rate and/or feed conversion in pigs, feedlot cattle, broilers and replacement pullets.

Store in a cool (5 – 15 °C) dry place.

Close bag after use.

**CONTAINS**

Each kilogram of premix contains 10 % m/m Tylosin phosphate.

**WARNINGS**

Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Withdraw Tyleco FG100 medicated feed from pigs 15 days before slaughter, and in broilers 3 days before slaughter.

**PRECAUTIONS AND SIDE EFFECTS**

Oedema of the anal mucose, partial anal protrusion, arythema and pruritus have occasionally been reported following tylosin medication in pigs.

Operators should wear protective clothin and masks when handling the product.

Wash hands thoroughly after use.

Do not use after expiry date indicated on the bag.

Medicated feed must be used within 3 months after mixing.

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

Thoroughly mix the required amount of TYLECO F.G. 100 according to the instructions below:

**Pigs**

<b>Pig mass (Feed category)</b>	<b>Gram TYLECO F.G. 100 premix per metric ton feed</b>	<b>Gram TYLOSIN per metric ton feed</b>
11 kg – 18 kg (Creep)	400	40
19 kg – 45 kg (Grower)	200 – 400	20 – 40
46 kg – market (Finisher)	200	20

It is recommended that TYLECO F.G. 100 is thoroughly mixed with a small quantity of finished feed (20 – 30 kg), before it is incorporated into the bulk of the feed being prepared. This procedure will assist in homogeneous distribution of TYLECO F.G. 100 in the final feed.

**Replacement Pullets**

In *Mycoplasma gallisepticum* possitive flocks – for improved feed efficiency.

Add 500 g TYLECO F.G. 100 per metric ton of complete feed to provide 50 ppm Tylosin.

In *Mycoplasma gallisepticum* negative flocks – for improved feed efficiency.  
Add 220 g TYLECO F.G. 100 per metric ton of complete feed to provide 22 ppm Tylosin.

**Broilers**

To improve the rate of mass gain and efficiency. Add 50 g TYLECO F.G. 100 per metric ton of complete feed to provide 5 ppm Tylosin.

**Feedlot cattle**

Add 100 g to 1 000 kg complete cattle feed to give 10 ppm Tylosin. Mix to provide 100 mg Tylosin per head per day.

There is no withdrawal period for TYLECO F.G. 100 in cattle.

**CONTENTS**

25 Kg

**Registration holder:**

ECO Animal Health Southern Africa (Pty) Ltd  
Co. Reg. No. 1992/00835/07  
P O Box 2009, FAERIE GLEN 0043, R.S.A.  
Tel: 012 991 6416

Lot No.

Exp.