

FOR ANIMAL USE ONLY

**TYLECO FG 250**  
Reg. No. G2956 Act/Wet 36/100

An aid in improving growth rate and/or feed conversion in pigs, feedlot cattle, broilers and replacement pullets.

**STORAGE INSTRUCTIONS**

Store below 25 °C in a dry place.

**COMPOSITION**

Each kilogram of premix contains 25 % m/m Tylosin phosphate.

**WARNINGS**

- WITHDRAW TYLECO 250 MEDICATED FEED FROM PIGS 15 DAYS BEFORE SLAUGHTER, AND IN BROILERS 3 DAYS BEFORE SLAUGHTER.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

**PRECAUTIONS AND SIDE EFFECTS**

- Do not use after expiry date indicated on the bag.
  - Medicated feed must be used within 3 months after mixing.
  - Operators should wear protective clothing and masks when handling the product.
  - Wash hands thoroughly after use.
- Oedema of the anal mucosa, erythema and pruritus have occasionally been reported following Tylosin medication in pigs.

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

Thoroughly mix the required amount of TYLECO FG 250 according to the instructions below:

**PIGS**

Body mass (kg)		g TYLOSIN FG 250 premix/metric ton feed	Concentration of Tylosin in final feed
11 – 18	Creep	160	40 ppm / dpm
19 – 45	Grower	80 – 160	20 – 40 ppm / dpm
46 - market	Finisher	80	20 ppm / dpm

It is recommended that TYLECO FG 250 is thoroughly mixed with a small quantity of finished feed (20 – 30 kg) before it is incorporated into the bulk of the feed being prepared. This procedure will assist in homogeneous distribution of TYLECO FG 250 in the final feed.

**REPLACEMENT PULLETS**

- *Mycoplasma gallisepticum* positive flocks: For improved feed efficiency add 200 g TYLECO FG 250 per metric ton of complete feed to provide 50 ppm Tylosin.
- *Mycoplasma gallisepticum* negative flocks: For improved feed efficiency add 88 g TYLECO FG 250 per metric ton of complete feed to provide 22 ppm Tylosin.

**BROILERS**

- To improve the rate of mass gain and efficiency: Add 20 g TYLECO FG 250 per metric ton of complete feed to provide 5 ppm Tylosin.

**FEEDLOT CATTLE**

- Add 40 g to 1 000 kg complete cattle feed to give 10 ppm Tylosin. Mix to provide 100 mg Tylosin per head per day.

**THERE IS NO WITHDRAWAL PERIOD FOR TYLECO FG 250 IN CATTLE.**

Presentation: 25 kg

**Registration holder/Registrasiehouer:**

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 1992/000835/07

P O Box 12406, Onderstepoort, 0110

Tel: (012) 545-0004