

**LABEL: ULTRATET L.A.
OUTER CONTAINER**

**FOR ANIMAL USE ONLY
SLEGS VIR DIEREGERBRUIK**

ULTRATET L.A.

Reg. No. G2857 Act/Wet 36/1947
N-SR0922

LONG ACTING INJECTABLE SOLUTION LANGWERKENDE INSPUITBARE OPLOSSING

For treatment of tick-borne-gallsickness (anaplasmosis), heartwater, foot-rot, pneumonia, navel-ill and joint-ill in livestock.
Vir die behandeling van bosluisoorgedraagde galsiekte (anaplasmosis), hartwater, vrotvootjie, longonsteking, naelstringsiekte en septiese gewrigsontsteking by vee.

STORE IN A COOL PLACE/BERG IN 'N KOEL PLEK

COMPOSITION/SAMESTELLING

Each 1 ml contains: Oxytetracycline hydrochloride 230 mg
Elke 1 ml bevat: Oksitetrasiklienhydrochloried 230 mg

40 ml	100 ml	500 ml	1 l
-------	--------	--------	-----

FOR FULL PARTICULARS SEE ENCLOSED PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUWILJET

Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd
ECO Dieregesondheid Suider-Afrika (Edms) Bpk
Co./Mpy. Reg. No. 92/00835/07
P O Box/Posbus 2009, FAERIE GLEN 0043
Tel: (012) 991 6416

Lot No.:	
Expiry/Verval:	

ECO Animal Health SA (Pty) Ltd
08/02/2010 Amendment to registration
1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)
2) Updated registration holder details

**LABEL: ULTRATET L.A.
IMMEDIATE CONTAINER**

**FOR ANIMAL USE ONLY
SLEGS VIR DIEREGERBRUIK**

ULTRATET L.A.

Reg. No. G2857 Act/Wet 36/1947
N-SR0922

LONG ACTING INJECTABLE SOLUTION LANGWERKENDE INSPUITBARE OPLOSSING

For treatment of tick-borne-gallsickness (anaplasmosis), heartwater, foot-rot, pneumonia, navel-ill and joint-ill in livestock.
Vir die behandeling van bosluisoorgedraagde galsiekte (anaplasrose), hartwater, vrotpootjie, longonsteking, naelstringsiekte en septiese gewrigsontsteking by vee.

STORE IN A COOL PLACE/BERG IN 'N KOEL PLEK

COMPOSITION/SAMESTELLING

Each 1 ml contains: Oxytetracycline hydrochloride 230 mg
Elke 1 ml bevat: Oksitetrasiklienhydrochloried 230 mg

40 ml	100 ml	500 ml	1 l
-------	--------	--------	-----

FOR FULL PARTICULARS SEE ENCLOSED PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET

Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd
ECO Dieregesondheid Suider-Afrika (Edms) Bpk
Co./Mpy. Reg. No. 92/00835/07
P O Box/Posbus 2009, FAERIE GLEN 0043
Tel: (012) 991 6416

Lot No.:	
Expiry/Verval:	

ECO Animal Health SA (Pty) Ltd

08/02/2010 Amendment to registration

1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)

2) Updated registration holder details

PACKAGE INSERT: ULTRATET L.A.

FOR ANIMAL USE ONLY

ULTRATET L.A.

Reg. No. G2857 Act 36/1947
N-SR0922

LONG ACTING INJECTABLE SOLUTION

STORAGE INSTRUCTIONS

Store in a cool place.

COMPOSITION

Each 1 ml contains Oxytetracycline hydrochloride 230 mg.

INDICATIONS

ULTRATET L.A. is indicated for the treatment and control of conditions caused by, or associated with oxytetracycline susceptible organisms. Its long action is recommended where for veterinary, practical or economic reasons it is not possible or desirable to handle and treat the animal frequently.

• **Cattle**

For the treatment of tick-borne gallsickness (anaplasmosis), heartwater, foot-rot, pneumonia, navel-ill and joint-ill.

• **Sheep and Goats**

For treatment of heartwater, foot-rot, pneumonia, navel-ill and joint-ill.

• **Pigs**

For treatment of foot-rot, pneumonia, navel-ill and joint-ill.

WARNINGS

Use only in cattle, sheep, goats and pigs.

DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 28 DAYS OF LAST TREATMENT.

DO NOT USE MILK FOR HUMAN CONSUMPTION WITHIN 5 DAYS OF LAST TREATMENT.

ECO Animal Health SA (Pty) Ltd

08/02/2010 Amendment to registration

1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)

2) Updated registration holder details

DO NOT ADMINISTER ULTRATET L.A. TO PIGLETS ON THE SAME DAY THAT THEY ARE INJECTED WITH IRON.

Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

Dosage

ULTRATET L.A. is designed for single intramuscular injection at 1 ml per 10 kg body mass which gives a dosage of 23 mg Oxytetracycline hydrochloride per kg body mass.

If the product is used intravenously, no long-acting effect will be achieved. Normally one injection is sufficient to treat diseases mentioned above. A second injection at 72 hours (3 days) after the initial injection may be given if necessary. Treatment should always be continued until total recovery.

For adult animals it is recommended that the dose be split and given in two injection sites.

Cattle: Not more than 20 ml per site.

Pigs: Not more than 10 ml per site.

Sheep and Goats: Not more than 5 ml per site.

For pigs under 10 kg it is preferable to inject subcutaneously. This route is not preferred for other animals as it may cause some swelling.

Administration

Injection in the neck is preferable to the rump as this tends to give higher antibiotic blood-levels and causes less tissue damage.

Inject deep into the fleshy part of the muscle. Temporary discomfort may be caused by intramuscular injection; massaging the site of injection will reduce discomfort and will help to disperse the antibiotic at the site of the injection.

N.B. for rapid recovery a prompt and accurate diagnosis is essential. It is, therefore, always advisable to consult your veterinarian for a correct diagnosis.

As with all oxytetracycline solutions darkening may occur during use without affecting efficacy.

PRESENTATION

40 ml, 100 ml, 500 ml and 1l polyethylene containers.

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 92/00835/07

P O Box 2009, FAERIE GLEN 0043

Tel: (012) 991 6416

ECO Animal Health SA (Pty) Ltd

08/02/2010 Amendment to registration

1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)

2) Updated registration holder details

VOUBILJET: ULTRATET L.A.

SLEGS VIR DIEREGERBUIK

ULTRATET L.A.

Reg. No. G2857 Wet 36/1947
N-SR0922

LANGWERKENDE INSPUITBARE OPLOSSING

BERGINGSAAWYSINGS

Berg in 'n koel plek.

SAMESTELLING

Elke 1 ml bevat Oksitetrasiklienhidrochloried 230 mg.

AANDUIDINGS

ULTRATET L.A. is vir die behandeling en beheer van siektes veroorsaak deur, of wat verband hou met organismes wat vatbaar is vir oksitetrasiklien. Die langwerking daarvan is aanbeveel waar vir veteriniere, praktiese of ekonomiese redes dit ontmoontlik of onwenslik is om diere meer dikwels te hanteer en te behandel.

- **Beeste**

Vir die behandeling van bosluisoorgedraagde galsiekte (anaplasmose), hartwater, vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking.

- **Skape en Bokke**

Vir die behandeling van hartwater, vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking.

- **Varke**

Vir die behandeling van vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking.

WAARSKUWINGS

Alleenlik vir gebruik by beeste, skape, bokke en varke.

MOENIE BEHANDELDE DIERE BINNE 28 DAE NA LAASTE BEHANDELING VIR MENSLIKE VERBRUIK SLAG NIE.

MOENIE MELK VIR MENSLIKE VERBRUIK BINNE 5 DAE NA LAASTE BEHANDELING GEBRUIK NIE.

MOENIE ULTRATET L.A. OP DIESELFDE DAG BY VARKIES SPUIT AS WAT HUL MET YSTER INGESPUIT IS NIE.

Hou buite bereik van kinders, oningeligte persone en diere.

ECO Animal Health SA (Pty) Ltd

08/02/2010 Amendment to registration

1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)

2) Updated registration holder details

Alhoewel hierdie middel breedvoerig onder 'n wye verskydenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

Dosis

ULTRATET L.A. is vir enkele binnespierse inspuiting ontwerp wat teen 1 ml per 10 kg liggaamsmassa 'n dosis van 23 mg Oksitetrasiklienhydrochloried per kg liggaamsmassa lewer.

As die produk binnears gebruik word sal geen langwerkende effek gehandhaaf word nie.

Normaalweg is een inspuiting voldoende om bogenoemde siektes te behandel. 'n Tweede inspuiting na 72 uur (3 dae) na die eerste inspuiting mag gegee word indien nodig.

Vir volwasse diere word dit aanbeveel dat die dosis verdeel word en op twee plekke ingespuut word.

Beeste: Nie meer as 20 ml per inspuitplek.

Varke: Nie meer as 10 ml per inspuitplek.

Skape en Bokke: Nie meer as 5 ml per inspuitplek.

Vir varke onder 10 kg is dit verkieslik om onderhuids in te spuit. Hierdie roete is nie verkieslik vir ander rasse nie omdat dit swelling kan veroorsaak.

Toediening

Inspuiting in die nek is verkieslik bo die kruis omdat dit geneig is om hoër antibiotikum bloedvlakke te gee en minder weefsel skade. Spuit diep in vleserige deel van die spier in. Tydelike ongemaklikheid na binnespierse inspuiting mag voorkom; masseering van die plek van inspuiting sal die ongemaklikheid verminder en help met die verspreiding van die antibiotikum by die plek van inspuiting.

L.W. Vir vinnige herstel is 'n spoedige en akkurate diagnose noodsaaklik. Dit is dus altyd raadsaam om u veearts vir 'n korrekte diagnose te raadpleeg.

Soos met alle oksitetrasiklien oplossings mag verdonkering gedurende gebruik voorkom sonder om die effektiwiteit te affekteer.

AANBIEDING

40 ml, 100 ml, 500 ml and 1l poliëtileen houers.

Registrasiehouer:

ECO Dieregesondheid Suider-Afrika (Edms) Bpk

Mpy. Reg. No. 92/00835/07

Posbus 2009, FAERIE GLEN 0043

Tel: (012) 991 6416

ECO Animal Health SA (Pty) Ltd

08/02/2010 Amendment to registration

1) Product name change from ECOMYCIN L.A. to ULTRATET L.A. (G2857)

2) Updated registration holder details