

Package Insert: ECOMECTIN PIG PREMIX

FOR ANIMAL USE ONLY

ECOMECTIN PIG PREMIX

Reg. No. G3860 Act 36/1947

CAUTION

STORAGE INSTRUCTIONS

Store in tightly closed, original container.

Store below 25 °C in a dry place.

Re-seal bag after use.

The product will remain stable in finished feed for 3 months when stored below 25 °C.

COMPOSITION

Ivermectin 0,6 % m/v

INDICATIONS

Antiparasitic remedy for pigs

For the treatment and control of gastro-intestinal roundworms, lungworms, lice and mange in adult and growing pigs.

- **Gastro-intestinal worms:**

Ascaris suum (adult and immature)

Hyostorngylus rubidus (adult and immature)

Oesophagostomum spp. (adult and immature)

Strongyloides ransomi (adult)

- **Lungworms:** *Metastrongylus* spp. (adult)

- **Lice:** *Haematopinus suis*

- **Mange mites:** *Sarcoptes scabiei var suis*

WARNINGS

- PIGS MUST NOT BE SLAUGHTERED FOR HUMAN CONSUMPTION DURING TREATMENT OR FOR 7 DAYS AFTER THE LAST TREATMENT.
- DO NOT USE FOR ANY OTHER ANIMAL SPECIES AS ADVERSE REACTIONS, INCLUDING FATALITIES IN DOGS, MAY OCCUR.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

SLEGS VIR DIEREGBRUIK

ECOMECTIN VOORMENGSEL VIR VARKE

Reg. No. G3860 Wet 36/1947

VERSIGTIG

BERGINGSAAWYSINGS

Berg in diggeseelde, oorspronklike houer.

Berg onder 25 °C in 'n droë plek.

Herseël houer na gebruik.

Die produk sal stabiel bly in die finale rantsoen vir 3 maande indien geberg onder 25 °C.

SAMESTELLING

Ivermectin 0,6 % m/v

INDIKASIES

Antiparasitiese middel vir varke

Vir die behandeling en beheer van rondewurms wat in die spysverteringskanaal voorkom, longwurms, luise en brandsiekte in volwasse en groeiende varke.

- **Wurms in die spysverteringskanaal:**

Ascaris suum (volwasse en onvolwasse)

Hyostorngylus rubidus (volwasse en onvolwasse)

Oesophagostomum spp. (volwasse en onvolwasse)

Strongyloides ransomi (volwasse)

- **Longwurms:** *Metastrongylus* spp. (volwasse)

- **Luise:** *Haematopinus suis*

- **Brandsiektemyte:** *Sarcoptes scabiei var suis*

WAARSKUWINGS

- VARKE MOENIE GEDURENDE BEHANDLING, OF VIR 7 DAE NA DIE LAASTE BEHANDLING VIR MENS-LIKE VERBRUIK GESLAG WORD NIE.
- MOENIE GEBRUIK IN ANDER DIERESPESIES NIE AANGESIEN ONGEWENSDE REAKSIES, INSLUITEND VREKTES BY HONDE, MAG VOORKOM.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

PRECAUTIONS

- Do not smoke or eat while handling the product.
- Wash hands after use.
- Use only in well ventilated areas or outdoors.
- In accidental skin contact occurs, wash the affected area immediately with soap and water.
- If accidental eye exposure occurs, flush eyes immediately with water and get medical attention.

Container disposal: Extremely dangerous to fish and aquatic life.

- Do not contaminate surface water or ditches with product or the used container.
- The container and residual contents should be disposed of safely by burying in waste ground away from water courses.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- Since the effect of ivermectin is not immediate, avoid contact between treated and untreated pigs for at least 1 week after completion of the treatment programme.
- Exposure of treated pigs to infected animals, contaminated premises, soil or pasture may result in reinfection.
- Since louse eggs can take up to 3 weeks to hatch, re-treatment may be necessary.

Dosage

- To ensure thorough dispersion of the product, it should first be mixed with a suitable quantity of feed ingredients before incorporation in the final mix.
- The product can be incorporated in pelleted feed preconditioned with steam for up to 10 seconds at a temperature not exceeding 85 °C.
- Ecomectin Pig Premix must be incorporated in feed and fed for 7 consecutive days at the recommended dose rate of 100 µg of ivermectin per kg of body mass per day.

Growing pigs

- Ecomectin Pig Premix must be thoroughly

VOORSORGMAATREËLS

- Moenie rook of eet terwyl die produk hanteer word nie.
- Was hande deeglik na gebruik.
- Gebruik slegs in areas met goeie ventillasie, of buitemuurs.
- Indien velkontak per ongeluk sou plaasvind, was die geaffekteerde area onmiddellik met seep en water.
- Indien die produk in die oë sou beland, spoel die oë dadelik met water uit en kry mediese hulp.

Verwydering van houer: Baie gevaarlik vir visse en ander waterlewe.

- Moenie oppervlak water of waterslote met die produk of leë houer besoedel nie.
- Die houer en oorblywende inhoud moet begrawe word in nuttelose grond weg van enige waterbron.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- Omrede ivermektien nie onmiddellik werk nie, voorkom kontak tussen behandelde en onbehandelde varke vir ten minste een week nadat die behandelingsprogram voltooi is.
- Behandelde varke wat blootgestel word aan besmette diere-, grond- of weiding, mag herbesmetting teweegbring.
- Omdat luis-eiers tot 3 weke lank kan neem om uit te broei, mag herbehandeling nodig wees.

Dosis

- Om deeglike verspreiding van die produk te verseker, moet dit eers met 'n geskikte kleiner hoeveelheid voer gemeng word voordat dat dit in die finale rantsoen ingesluit word.
- Die produk kan in verkorrelde voer ingesluit word, waar die voer met stoom voorberei word vir 10 sekondes teen 'n temperatuur nie hoër as 85 °C nie.
- Ecomectin Voormengsel vir varke moet by die voer ingemeng word en vir 7 agtereenvolgende dae teen die voorgestelde dosis van 100 µg ivermektien per liggaamsmassa per dag gevoer word.

Groeinde varke

- Ecomectin Voormengsel vir varke moet

mixed in the finished feed and fed continuously as the only ration for 7 consecutive days.

- For pigs up to 40 kg body mass, include 333 g of Ecomectin Pig Premix in each tonne of finished feed.
- For pigs weighing 40 – 100 kg, include 400 g of Ecomectin Pig Premix per tonne of finished feed.

Adult pigs

- The recommended dose level for adult pigs is 1,67 kg of Ecomectin Pig Premix in each tonne of finished feed.
- The resulting medicated feed should be fed at a rate of 1 kg/100 kg body mass each day for 7 days as part of the ration.

- Where medicated feed is fed as part of the ration it is recommended that the Ecomectin Pig Premix medicated feed is fed first.
- After this is consumed, any remaining daily feed allocation should be given.
- This should be repeated for 7 consecutive days.
- Alternatively, where dry feed intake can be accurately measured and all the animals to be treated have a similar body mass, the inclusion rate can be calculated using the following formula. This assumes the total ration is to be medicated.

$$\text{Premix inclusion rate (kg/tonne of feed)} = \frac{0,1 \times \text{average body mass (kg)}}{6 \times \text{average daily feed intake (kg)}}$$

Treatment programme

Growing pigs

- Groups of growing pigs should be treated for 7 consecutive days on transfer to clean quarters.
- Where an all-in-all-out system is not possible, it is recommended that the parasite control programme should start with all the growing pigs already in the pen.

Adult pigs

- All breeding pigs should be treated in a blanket programme every 6 months or as often as necessary.
- A treatment will be of particular use

deeglik in die finale rantsoen ingemeng word en aanhoudend gevoer word as die enigste rantsoen vir 7 agtereenvolgende dae.

- Vir varke tot en met 40 kg liggaamsmassa, sluit 333 g Ecomectin Voormengsel in elke ton finale rantsoen in.
- Vir varke vanaf 40 – 100 kg, sluit 400 g Ecomectin Voormengsel in per ton van die finale rantsoen.

Volwasse varke

- Die aanbevole dosis is 1,67 kg Ecomectin Voormengsel per ton finale rantsoen.
- Die gemedikeerde rantsoen moet gevoer word teen 1 kg pr 100 kg liggaamsmassa elke dag vir 7 dae as gedeelte van die totale rantsoen.

- Indien gemedikeerde voer as deel van die rantsoen gevoer word, word aanbeveel dat die gemedikeerde voer eerste gevoer word.
 - Nadat die gemedikeerde voer opgevreet is, kan die oorblywende gedeelte van die daaglikse rantsoen gevoer word.
 - Dit moet herhaal word vir 7 agtereenvolgende dae.
 - Andersins, indien droë voer akkuraat gemeet kan word en alle diere wat behandel moet word min of meer ewe veel weeg, kan die insluitingsvlakke bereken word deur die volgende formule te gebruik. Die aanname word gemaak dat die totale rantsoen gemedikeer is.
- $$\text{Voormengsel insluitings-tempo (kg/ton voer)} = \frac{0,1 \times \text{gemiddelde liggaamsmassa (kg)}}{6 \times \text{gemiddelde daaglikse voerinnome (kg)}}$$

Behandelings-program

Groeierende varke

- Groepe groeiende varke moet vir 7 agtereenvolgende dae behandel word wanneer hulle na skoon hokke verskuif word.
- Waar 'n alles-in, alles-uit sisteem nie moontlik is nie, word voorgestel dat die parasietbeheer program moet begin by die groeiende varke wat reeds in die hok is.

Volwasse varke

- Kombers-behandeling moet elke 6 maande plaasvind in alle teelvarke. Andersins, so dikwels as wat nodig is.
- 'n Behandeling sal van goeie waarde wees

before moving sows to clean quarters or a fresh paddock.

Pregnancy/lactation

- Ecomectin Pig Premix can be administered to sows at any stage of pregnancy or lactation.
- Ecomectin Pig Premix given to pregnant sows before farrowing effectively controls transmission via the milk of *S. ransomi* to piglets.
- Ecomectin Pig Premix can be used in breeding sows and boars and will not affect fertility.

PRESENTATION

333 g, 5 kg

REGISTRATION HOLDER:

ECO Animal Health Southern Africa (Pty) Ltd
Co. reg. no. 1992/00835/07
P O Box 2009, FAERIE GLEN
0043, RSA
Tel: +27 12 817 9060
Helpline: 0860 833 2787

DISTRIBUTED BY:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009, FAERIE GLEN
0043, RSA
Tel: +27 12 817 9060
Helpline: 0860 833 2787
www.afrivet.co.za

net voor soê na 'n skoon huis verskuif word.

Dragtigheid/laktasie

- Ecomectin Voormengsel kan op enige stadium van dragtigheid of laktasie aan sôe toegedien word.
- Ecomectin Voormengsel vir varke wat gegee word aan dragtige sôe voor jong, beheer die oordrag van *S. ransomi* deur die melk na varkies doeltreffend.
- Ecomectin Voormengsel kan ook gebruik word in teelsôe- en bere en sal nie vrugbaarheid beïnvloed nie.

AANBIEDING

333 g, 5 kg

REGISTRASIEHOUER:

ECO Dieregesondheid Suider Afrika (Edms) Bpk
Mpy. Reg. no. 1992/00835/07
Posbus 2009, FAERIE GLEN
0043, RSA
Tel: +27 12 817 9060
Hulplyn: 0860 833 2787

VERSPREI DEUR:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009, FAERIE GLEN
0043, RSA
Tel: +27 12 817 9060
Hulplyn: 0860 833 2787
www.afrivet.co.za