

## **PACKAGE INSERT: ECOSULF L.A.**

### **ECOSULF L.A.**

RSA Reg. No. G3037 Act 36/1947

Namibia S0 V02/17.1.7/705

### **STORAGE INSTRUCTIONS**

Store in a cool, dark place.

### **COMPOSITION**

Contains Sodium sulphadimethoxine 40 % m/v.

### **INDICATIONS**

For the treatment of footrot, navel-ill, joint-ill and pneumonia in stock, coccidiosis and bacterial scours in calves and lambs and strangles in horses.

### **WARNINGS**

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- MILK FROM TREATED COWS MUST NOT BE USED FOR HUMAN CONSUMPTION WITHIN 48 HOURS OF LAST TREATMENT.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### **DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

#### **Administration**

Administer by subcutaneous or intravenous routes.

Massage the injection site to maximise absorption and to minimise swelling.

When using the intravenous route, give the injection very slowly.

Discontinue for a while at the first sign of discomfort and continue only when all seems normal.

**N.B.** Treated animals must have free access to ample drinking water during treatment and for at least two days after the last treatment.

#### **Indications**

- **Cattle, sheep and goats:**  
Footrot, pneumonia, navel-ill and joint-ill.
- **Horses:**  
Pneumonia, strangles, joint-ill and navel-ill.
- **Pigs:**  
Pneumonia and joint-ill.
- **Calves and lambs:**  
Coccidiosis and bacterial scours.

**Dosage**

An initial dosage double that of the maintenance dosage is recommended.

**AFTER ADMINISTERING THE INITIAL DOSE, ADMINISTER THE MAINTENANCE DOSE ONCE A DAY FOR 3 DAYS.**

DO NOT TREAT FOR MORE THAN 4 DAYS.

<b>Species</b>	<b>Initial dose</b>	<b>Maintenance dose</b>
Cattle, horses, sheep, goats and pigs	1 – 2 ml / 10 kg body mass	0,5 – 1 ml / 10 kg body mass

**N.B.** Consult your veterinarian immediately if any uncertainty arises as to diagnosis or dosage or if little or no improvement is effected by the treatment.

**Packaging Size:**

100 ml, 200 ml, 250 ml, 500 ml

**Registration holder/Registrasiehouer:**

ECO Animal Health Southern Africa (Pty) Ltd  
Co./Mpy. Reg. No. 1992/000835/07  
P O Box/Posbus 2009, Faerie Glen  
0043, R.S.A.  
Helpline/Hulplyn: 0860 833 2787  
[www.ecoanimalhealth.com](http://www.ecoanimalhealth.com)

**Distributed by/Versprei deur:**

Afrivet Business Management (Pty) Ltd  
Co./Mpy. Reg 2000/011263/07  
PO Box/Posbus 2009, Faerie Glen  
0043, RSA  
Helpline/Hulplyn: 0860 833 2787  
[www.afrivet.co.za](http://www.afrivet.co.za)

## **VOUBILJET: ECOSULF L.A.**

### **ECOSULF L.A.**

RSA Reg. No. G3037 Wet 36/1947

Namibia S0 V02/17.1.7/705

### **BERGINGSAAWYSINGS**

Berg in 'n koel, donker plek.

### **SAMESTELLING**

Bevat Natriumsulfadimetoksien 40 % m/v.

### **INDIKASIES**

Vir die behandeling van vrotpootjie, naelstringsiekte, septiese gewrigsontsteking en longontsteking by vee, koksidiöse en bakteriese maagwerking by kalwers en lammers en nuwesiekte by perde.

### **WAARSKUWINGS**

- MOENIE DIERE BINNE 7 DAE NA DIE LAASTE BEHANDELING VIR MENSLIKE VERBRUIK SLAG NIE.
- MELK VAN BEHANDELDE KOEIE MOET NIE VIR MENSLIKE VERBRUIK BINNE 48 UUR NA LAASTE BEHANDELING GEBRUIK WORD NIE.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

### **GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI**

#### **Toediening**

Dien toe deur onderhuidse- of binnearse inspuiting.

Masseer die inspuitplek om opname te bevorder en swelling te verminder.

Die binnearse inspuiting moet baie stadig toegedien word.

Onderbreek die inspuiting by die eerste teken van ongemak en hervat inspuiting wanneer die dier normal voorkom.

**L.W.** Behandelde diere moet vrye toegang tot volop drinkwater hê tydens behandeling en minstens twee dae na die laaste toediening.

#### **Aanduidings**

- **Beeste, skape en bokke:**  
Vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking.
- **Perde:**  
Longontsteking, nuwesiekte, septiese gewrigsontsteking en naelstringsiekte.
- **Varke:**  
Longontsteking en septiese gewrigsontsteking.
- **Kalwers en lammers:**  
Koksidiöse en bakteriese maagwerking.

## Dosis

'n Aanvangsdosis dubbel die hoeveelheid van die onderhoudsdosis word aanbeveel.

**NADAT DIE AANVANKLIKE DOSIS TOEGEDIEN IS, DIEN DIE ONDERHOUDSDOSIS EEN MAAL 'N DAG VIR 3 DAE TOE.**

MOET NIE VIR LANGER AS 4 DAE BEHANDEL NIE.

Diersoort	Aanvangsdosis	Onderhoudsdosis
Beeste, perde, skape, bokke en varke	1 – 2 ml / 10 kg liggaamsmassa	0,5 – 1 ml / 10 kg liggaamsmassa

**L.W.** Indien enige onsekerheid ten opsigte van die dosis of diagnose ontstaan, of as geen beterskap intree na die behandeling nie, moet u onmiddellik u veearts raadpleeg.

## AANBIEDING:

100 ml, 200 ml, 250 ml, 500 ml

### Registration holder/Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd  
Co./Mpy. Reg. No. 1992/000835/07  
P O Box/Posbus 2009, Faerie Glen  
0043, R.S.A.  
Helpline/Hulplyn: 0860 833 2787  
[www.ecoanimalhealth.com](http://www.ecoanimalhealth.com)

### Distributed by/Versprei deur:

Afrivet Business Management (Pty) Ltd  
Co./Mpy. Reg 2000/011263/07  
PO Box/Posbus 2009, Faerie Glen  
0043, RSA  
Helpline/Hulplyn: 0860 833 2787  
[www.afrivet.co.za](http://www.afrivet.co.za)