



CLASS III

CAUTION
VERSIGTIG

Blue band

ERADIFLY GRANULAR

L10328 Act 36 of 1947

Granule

ACTIVE INGREDIENT

Novaluron 0,2 % (IGR -insect growth regulator)

PRODUCT INFORMATION

ERADIFLY GRANULAR is a larvicide with specific properties developed for the control of flies in manure and other breeding sites. ERADIFLY GRANULAR provides effective control in cattle operations. Fly control in cattle operations should include appropriate sanitary and management practices to reduce the number of size of insect breeding sites.

ERADIFLY GRANULAR affects only developing immature stages by disrupting the molting process. Consequently, fully developed adult stage of pest and beneficial species are not affected. The elimination of adult insects occurs gradually and become visible one to two weeks after application.

WARNINGS

- Prevent contamination of feed or water troughs.
- Do not apply directly ERADIFLY GRANULAR to livestock or livestock feed as illegal residues may result.
- Do not feed manure treated with ERADIFLY GRANULAR to animals.
- Causes moderate eye irritation.
- Harmful if absorbed through the skin.
- Avoid contact with skin, eyes or clothing.
- Wear protective gloves during use.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

ERADIFLY GRANULAR

L10328 Wet 36 of 1947

Korreltjie

AKTIEWE BESTANDDEEL

Novaluron 0,2 % (IGR - insek groeireguleerde)

PRODUKINLIGTING

ERADIFLY KORRELS is 'n larwedoder met spesifieke eienskappe, ontwikkel vir die beheer van vlieë in mis en ander broeiplekke. ERADIFLY KORRELS bied doeltreffende beheer in veebedrywighede. Vliegbeheer in veebedrywighede moet toepaslike sanitêre en bestuurspraktyke insluit om die aantal en grootte van insekbroeiplekke te verminder.

ERADIFLY KORRELS affekteer net die ontwikkeling van onvolwasse stadiums deur die versteuring van die vervellingsproses . Gevolglik word ten volle ontwikkelde volwasse stadium van plaag-en voordelige spesies nie geraak nie. Die uitskakeling van volwasse insekte gebeur geleidelik en is sigbaar een tot twee weke na toediening.

WAARSKUWINGS

- Voorkom besoedeling van voedsel of waterkrippe.
- Moenie ERADIFLY GRANULAR direk op diere of dierevoer toedien nie. Dit mag reste in eetbare weefsel veroorsaak.
- Moenie mis wat met ERADIFLY GRANULAR behandel is, aan diere voer nie.
- Veroorsaak matige oog irritasie .
- Skadelik indien deur die vel geabsorbeer word.
- Vermy kontak met die vel , oë of klere.
- Dra beskermende handskoene tydens gebruik.
- Was deeglik met seep en water na hantering en voordat jy eet , drink , kougom kou of tabak gebruik.
- Hou buite bereik van kinders, oningelege persone en diere.
- Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, mag dit faal as gevolg van 'n wye verskeidenheid van redes. Indien dit vermoed word , raadpleeg 'n veearts en verwittig die registrasiehouer .



FIRST AID

In the event of any adverse reactions contact a doctor and make this label available.

Eyes

- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Skin

- Remove contaminated clothing.
- Rinse skin immediately with plenty of water for 15 – 20 minutes.

Ingestion

- Contact a doctor immediately for treatment advice.
- Do not give any liquid to the person.
- Do not induce vomiting unless told to do so by a doctor.
- Do not give anything by mouth to an unconscious person.

Inhalation

- Move person to fresh air.
- If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to mouth if possible.
- Consult a doctor for further treatment advice.

PRECAUTIONS

- This pesticide is toxic to freshwater and estuarine/marine invertebrates. Runoff can be hazardous to aquatic organisms in water adjacent to treated areas.
- This product is intended for use only in sites listed on this label.
- Do not apply to intertidal area below the mean high water mark.
- Do not contaminate water when disposing of equipment, wash waters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place. Do not store this product under wet conditions. Avoid sunlight and temperature exceeding 35 °C.

Pesticide disposal: Waste resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container disposal: Do not reuse or refill this container. Completely empty contents before disposal of the container. Puncture and dispose of

NOODHULP

In die geval van enige nadelige reaksies kontak 'n geneesheer en maak hierdie etiket beskikbaar.

Oë

- Hou oog oop en spoel stadig en versigtig met water vir 15 - 20 minute.
- Verwyder kontaklense, indien teenwoordig, na die eerste 5 minute, en gaan voort om oog te spoel.

Vel

- Verwyder besoedelde klere.
- Spoel vel dadelik met baie water vir 15 - 20 minute

Inname

- Kontak 'n dokter onmiddellik vir behandelingsadvies.
- Moenie enige vloeistof vir die persoon gee nie.
- Moenie braking veroorsaak nie, tensy dit deur 'n dokter aanbeveel word.
- Moet niks per mond aan 'n bewusteloze persoon gee nie.

Inaseming

- Neem persoon na vars lug.
- Indien die persoon nie asemhaal nie, bel 'n ambulans en pas dan kunsmatige asemhaling toe, verkieslik mond-tot-mond indien moontlik.
- Spreek 'n dokter vir verdere behandelingsadvies.

VOORSORGMAATREËLS

- Hierdie plaagdoder is giftig vir varswater- en estuariese/mariene ongewerweldes. Afloopwater kan gevaarlik wees vir waterorganismes in die water langs behandelde areas.
- Hierdie produk is bedoel slegs vir gebruik in plekke op die etiket gelys.
- Moenie toedien op intergety gebied onder die gemiddelde hoogwaterlyn nie.
- Moenie water besoedel met die wegdoen van houers, waswater of afspoelwater nie.

STOOR EN WEGDOENNING

Moenie water, voedsel of voer besoedel deur die stoor of wegdoenning nie.

Storage: Berg in 'n koel, droë plek. Moet nie hierdie produk stoor in nat toestande nie. Vermy sonlig en temperatuur van meer as 35 °C.

Plaagdoder wegdoenning: Afval wat voortspruit uit die gebruik van hierdie produk, moet weggedoen word op die terrein of by 'n goedgekeurde stortingsfasiliteit.

Houervernietiging: Moenie hierdie houer hergebruik of hervul nie. Maak houer heeltemal

CAUTION
VERSIGTIG

CLASS III

in a sanitary landfill or by incineration. If burned, stay clear of smoke.

Blue band

leeg voor vernietiging. Steek vol gate en gooi weg in 'n sanitêre stortingsterrein of vernietig deur dit te verbrand. As verbrand word, bly weg van die rook .

DIRECTIONS FOR USE: Use only as directed

ERADIFLY GRANULAR is designed for dry scattering. It can be used alternatively with fly adulticides and is an important component of any integrated fly control programme.

Application sites

Apply only on fly breeding sites. It is not intended for application to fly resting sites (walls, ceilings, etc.), as it has no effect on adult flies.

Application method

Dry scattering – apply directly and evenly on the insect breeding sites. Wear gloves when scattering.

Application schedule

The recommended treatment interval depends on management and housing systems as well as on climatic conditions. More specific recommendations are given in the application schedules below.

Cattle Operations

Apply NOVALURON 0.2G directly to fly breeding sites around round hay bales, dairy barns, cattle feedlots, and all other premise areas where flies can breed. Apply 26 g/m².

Apply NOVALURON 0.2G approximately 3 days after manure removal and repeat treatment after each removal or when fly larvae are found.

Application rate

Scatter granules at a rate of 26 g per m² of soil. Best results can be expected in wet soil, with enough animals present to trample the granules into the soil.

Packaging Size:

4,5 Kg, 22,5 Kg, 45 Kg

Registration holder:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Helpline: 0860 833 2787

UN No. 3082

GEBRUIKSAANWYSINGS : Gebruik slegs soos aangedui

ERADIFLY KORRELS is ontwerp vir droë uitstrooiing. Dit kan alternatiewelik met volwasse vlieg doders gebruik word en is 'n belangrike komponent van enige geïntegreerde vliegbeheer program.

Aanwendingsplekke

Slegs van toepassing op vliegbroeiplekke. Dit is nie bedoel vir die aanwending op vlieg rusplekke (mure, plafonne, ens) nie, want dit het geen invloed op volwasse vlieë nie.

Aanwendings metode

Droë uitstrooiing - regstreeks en egalig op die insek broeiplekke. Dra handskoene wanneer uitstrooi.

Aanwendings skedule

Die aanbevole behandelingsinterval is afhanklik van die bestuur en behuisingsstelsel, sowel as klimaatstoestande. Meer spesifieke aanbevelings word in die aanwendingskledules hieronder gegee.

Vee bedrywighede

Strooi NOVALURON 0.2G direk op vlieg broeiplekke rondom bale hooi, melkskure, beesvoerkrale, en alle ander areas waar vlieë kan teel. Strooi 26 g/m². Strooi NOVALURON 0.2G ongeveer 3 dae nadat die mis verwyder is en herhaal die behandeling na elke verwydering of wanneer vlieg larwes gevind word..

Aanwendingsdigtheid

Strooi die korrels teen 'n digtheid van 26 g per m² grond. Die beste resultate kan verwag word in nat grond, met genoeg diere teenwoordig om die korrels in die grond in te trap.

Aanbieding:

4,5 Kg, 22,5 Kg, 45 Kg

Registrasiehouer:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787

VN Nr. 3082