

**PACKAGE INSERT: Eraditick® 125**

FOR EXTERNAL ANIMAL USE ONLY

**ERADITICK® 125**

RSA Reg. No. G3583 Act 36/1947

Namibia  V10/18.3.4/1147

Botswana W1301247

**Emulsifiable concentrate**

**OXPECKER COMPATIBLE**

**HARMFUL**

**STORAGE INSTRUCTIONS**

Protect product from extremes of heat and cold and from dampness.

**COMPOSITION**

Amitraz 12,5 % m/v

**INDICATIONS**

CATTLE: Controls ticks. Kills lice and mange mites.

SHEEP AND GOATS: Controls ticks. Controls sheep scab. Kills itch mites and goat mange mites.

**WARNINGS**

- **WITHDRAWAL PERIOD:**
- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST DIPPING.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this label available to him.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

**PRECAUTIONS**

- Tightly reseal the container if all the contents have not been used.
- Destroy empty containers.
- Wash hands thoroughly after use.

**DIRECTIONS FOR USE – USE ONLY AS DIRECTED**

For external use on cattle, sheep and goats.

ERADITICK® 125 should not be mixed with any other dip.

**CATTLE**

• **Spraying**

Dipwash should be freshly prepared for each spraying.

Add required amount of ERADITICK® 125 to ± 10 ℓ water in a suitable container.

When well mixed add to measured bulk water in spray sump and stir thoroughly.

Use at a rate as per the table below:

<b>ERADITICK® 125</b>	<b>WATER</b>
100 mℓ	50 ℓ
200 mℓ	100 ℓ
250 mℓ	125 ℓ
500 mℓ	250 ℓ
1 ℓ	500 ℓ
5 ℓ	2 500 ℓ
10 ℓ	5 000 ℓ
20 ℓ	10 000 ℓ

## Boost

It is recommended that not more than 400 head be sprayed in the spray race at one time, especially where fouling is a problem. If, however, this figure is exceeded, boosting at the rate of 100 mℓ per 100 head, commencing at 400 head, can be undertaken to maintain a satisfactory deposit of active ingredient, especially where tick-borne diseases are a problem.

### • Plunge dip

**Total replenishment:** This requires that the diptank is topped up to the desired level on each dipping occasion. ERADITICK® 125 is then added on every occasion at a rate of 1 ℓ to 2 500 ℓ water, taking the total amount of water in the tank into the calculation.

### Lime stabilised:

- **Fresh fill:** Add ERADITICK® 125 at the rate of 1 ℓ per 500 ℓ water.

At the end of each dipping add 5 kg calcium hydroxide (75 %) per 100 head of cattle dipped.

- **Replenishment:** Add ERADITICK® 125 at the rate of 1 ℓ per 500 ℓ water added.

At the end of each dipping, calcium hydroxide (75 %) must be added to the tank at a rate of 5 kg per 100 head of cattle dipped to ensure that the dipwash pH is in the region of 12.

### • “Head Count” Replenishment System

After dipping add water to the diptank to fill the diptank to the required level. Use the following table to calculate how much ERADITICK® 125 needs to be added to the diptank.

Number of animals dipped	Quantity of ERADITICK® 125 to be added to the diptank
10	55.6 mℓ
20	111.2 mℓ
50	278 mℓ
100	556 mℓ
1000	5.56 ℓ

Note: 1ℓ of ERADITICK® 125 will replenish sufficient dipwash to dip 179 cattle.

At the end of each dipping, calcium hydroxide (75 %) must be added to the tank at a rate of 5 kg per 100 head of cattle dipped to ensure that the dipwash pH is in the region of 12.

### Frequency:

- **Ticks and mange** Spray/dip weekly
- **Lice** Spray/dip twice with a 7 day interval

Should dipping intervals extend beyond 3 weeks, additional lime should be added at a rate of 1 kg lime (75 % active) per 1 000 ℓ water in the diptank. The dipwash should be adequately protected for 2 months if this procedure is followed.

## SHEEP AND GOATS

ERADITICK® 125 should be administered by means of either:

1. 2 Dippings using fresh, unstabilised dipwash. ERADITICK® 125 should be added at a rate of 1 ℓ per 375 ℓ water, taking the total amount of water in the tank into consideration.  
OR
2. A fresh, unstabilised dipwash for the first dipping, which is then stabilised with the addition of lime at a rate of 4 kg lime per 750 ℓ dipwash, added immediately after the initial dipping. This stabilised dipwash is then suitable for the second dipping, 7 days later.

### Frequency:

Sheep and goats should be dipped twice with a 7 day interval to kill sheep scab, itch and goat mange mites. Ensure that the animals are immersed for approximately 1 minute with the heads being submerged at least 3 times in order to ensure that the inside of the ears are thoroughly wetted.

**N.B.** Sheep scab is a notifiable disease and must be reported to the nearest State Veterinarian.

### Presentation

100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ, 1 ℓ, 5 ℓ, 10 ℓ, 20 ℓ

**Registration holder:**

Afrivet Business Management (Pty) Ltd

Co. Reg 2000/011263/07

P O Box 2009, Faerie Glen

0043, Pretoria, R.S.A.

Helpline: 0860 833 2787

[www.afrivet.co.za](http://www.afrivet.co.za)

SLEGS VIR UITWENDIGE DIERGEBRUIK

## ERADITICK® 125

RSA Reg. No. G3583 Wet 36/1947

Namibia SO V10/18.3.4/1147

Botswana W1301247

**Emulsifiseerbare konsentraat**

**RENOSTERVOËL VERSOENBAAR**

**SKADELIK**

**BERGINGSAAWYSINGS**

Beskerm teen uitermatige hitte en koue en teen klamheid.

**SAMESTELLING**

Amitras 12,5 % m/v

**INDIKASIES**

BEESTE: Beheer bosluise. Dood luise en skurfte myte.

SKAPE EN BOKKE: Beheer bosluise. Beheer skaapbrandsiekte. Dood jeukmyte en bokskurfte myte.

**WAARSKUWINGS**

- **ONTREKKINGSPERIODE:**
- MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 7 DAE VANAF DIE LAASTE DIPPING SLAG NIE.
- MOENIE OP PERDE GEBRUIK NIE.
- In geval van vergiftiging raadpleeg 'n geneesheer en stel hierdie etiket aan hom beskikbaar.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

**VOORSORGMATREËLS**

- Herseël die houer dig indien al die inhoud nie gebruik is nie.
- Vernietig leë houers.
- Was hande deeglik na gebruik.

**GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI**

Vir uitwendige gebruik op beeste, skape en bokke.

ERADITICK® 125 behoort nie met enige ander dip gemeng te word nie

**BEESTE**

- **Bespuiting**

Dipmengsel behoort vars opgemaak te word voor elke bespuiting.

Voeg die vereiste hoeveelheid ERADITICK® 125 by ± 10 ℓ water in 'n geskikte houer.

Wanneer dit deeglik gemeng is voeg dit by afgemete volume water in die dipmengselhouer en roer deeglik.

Gebruik teen 'n koers soos in die tabel hieronder:

ERADITICK® 125	WATER
100 mℓ	50 l
200 mℓ	100 l
250 mℓ	125 ℓ
500 mℓ	250 l
1 ℓ	500 l
5 ℓ	2 500 l
10 ℓ	5 000 l
20 ℓ	10 000 l

### Versterking

Dit word aanbeveel dat nie meer as 400 beeste op een tydstep in die spuitdip bespuit word nie, veral waar besoedeling 'n probleem is. As hierdie getal oorskrei word kan versterking teen 'n koers van 100 mℓ per 100 diere, met aanvang by 400 diere, gedoen word om 'n aanvaarbare neerslag van aktiewe bestanddeel te verseker, veral waar bosluis oordraagbare siektes 'n probleem is.

#### • Dompeldip

**Totale aanvulling:** Dit verg dat die dipbak met elke dipping tot by die gewenste vlak aangevul word. ERADITICK® 125 word dan met elke dipping bygevoeg teen 'n koers van 1 ℓ by 2 500 ℓ water, met die totale hoeveelheid water in die dipbak in berekening gebring.

#### Stabilisering met kalk:

- **Varsvulling** Voeg ERADITICK® 125 by teen 'n koers van 1 ℓ per 500 ℓ water.  
5 kg Kalsiumhidroksied (75 %) per 100 kop beeste gedip moet na dipping by die dipmengsel bygevoeg word.
- **Aanvulling** Voeg ERADITICK® 125 by teen 'n koers van 1 ℓ per 500 ℓ water wat bygevoeg is.  
Aan die einde van elke dipping, moet kalsiumhidroksied (75 %) teen 'n koers van 5 kg per 100 kop beeste gedip by die dipbak gevoeg word, om te verseker dat die dipmengsel se pH in die omgewing van 12 is.

#### • "Head Count" Aanvulling Stelsel

Na dipping, voeg water by die dipbak tot by die nodige vlak. Gebruik dan die onderstaande tabel om te bereken hoeveel ERADITICK® 125 nodig is vir aanvulling.

Aantal diere gedip	Hoeveelheid ERADITICK® 125 wat by die dipbak gevoeg moet word
10	55.6 mℓ
20	111.2 mℓ
50	278 mℓ
100	556 mℓ
1000	5.56 ℓ

Nota: 1 ℓ ERADITICK® 125 sal genoeg dipmengsel aanvul vir die dip van 179 beeste.

Aan die einde van elke dipping, moet kalsiumhidroksied (75 %) teen 'n koers van 5 kg per 100 kop beeste gedip by die dipbak gevoeg word, om te verseker dat die dipmengsel se pH in die omgewing van 12 is.

#### Diptussenpose:

- **Bosluis en myte** Bespuit/dip weekliks
- **Luise** Bespuit/dip twee maal met 'n tussenpose van 7 dae

As die dip tussenpose langer as 3 weke is, moet addisionele kalk bygevoeg word teen 'n koers van 1 kg kalk (75 % aktief) per 1 000 ℓ water in die dip. As hierdie prosedure gevolg word behoort dit die dip vir 2 maande voldoende te beskerm.

### SKAPE EN BOKKE

ERADITICK® 125 behoort soos volg aangewend te word:

1. 2 Dippings met vars, ongestabiliseerde dipmengsel. ERADITICK® 125 moet bygevoeg word teen 'n koers van 1 ℓ per 375 ℓ water, met die totale hoeveelheid water in die dipbak in berekening gebring.  
OF
2. 'n Vars, ongestabiliseerde dipmengsel vir die eerste dipping, wat dan gestabiliseer word met die byvoeging van kalk, teen 'n koers van 4 kg kalk per 750 ℓ dipmengsel, wat onmiddellik na die eerste dipping bygevoeg word. Hierdie gestabiliseerde dipmengsel is dan geskik vir die tweede dipping 7 dae later.

#### Diptussenpose:

Skaap en bokke moet twee keer gedip word met 'n 7 dae tussenpose om skaapbrandsiekte-, jeuk- en bokskurfte myte te dood. Verseker dat die diere vir ongeveer 1 minuut gedompel is, en dat hulle koppe ten minste 3 keer onderdompel word, om deeglike benatting van die binnekant van die ore te verseker.

**L.W.** Skaapbrandsiekte is 'n aanmeldbare siekte en moet by die naaste Staatsveearts aangemeld word.

**Aanbieding**

100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ, 1 ℓ, 5 ℓ, 10 ℓ, 20 ℓ

**Registrasiehouer:**

Afrivet Business Management (Pty) Ltd

Mpy. Reg 2000/011263/07

Posbus 2009, Faerie Glen

0043, Pretoria, R.S.A.

Hulplyn: 0860 833 2787

[www.afrivet.co.za](http://www.afrivet.co.za)