

Package Insert: Eraditick® Cattle Pour-On

FOR EXTERNAL ANIMAL USE ONLY

ERADITICK® CATTLE POUR-ON

RSA Reg. No. G4254 Act 36/1947
Namibia  V04/18.3.4/540

HARMFUL

COMPOSITION

Amitraz 2,0 % m/v

INDICATIONS

READY-TO- USE POUR-ON FOR THE CONTROL OF TICKS ON CATTLE

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness. Store away from food and feed. Tightly reseal the container if all the contents have not been used.

WARNINGS

- DO NOT USE ON HORSES.
- DO NOT USE ON CALVES UNDER 1 MONTH.
- Withdrawal period:
 - Milk: 2 days
 - Meat: 7 days
- Signs of salivation and lethargy may occur with overdosing. These signs are transient and will pass with no ill effect.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.
- **N.B.** Pyrethroid pour-ons may cause irritation. Treat a small group of animals first and observe for an hour before treating the rest of the herd.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- FOR EXTERNAL USE ON CATTLE ONLY

SLEGS VIR UITWENDIGE DIEREGERBUIK

ERADITICK® BEES OPGIETMIDDEL

RSA Reg. No. G4254 Act 36/1947
Namibia  V04/18.3.4/540

SKADELIK

SAMESTELLING

Amitras 2,0 % m/v

INDIKASIES

GEREED-VIR-GEbruik OPGIETMIDDEL VIR DIE BEHEER VAN BOSLUISE OP BEESTE

BERGINGSAAWYSINGS

Beskerm teen uiterstes van hitte, koue en teen klammigheid. Berg weg van voedsel en voer. Indien inhoud nie opgebruik word nie, herseël houër deeglik.

WAARSKUWINGS

- MOENIE OP PERDE GEBRUIK NIE.
- MOENIE OP KALWERS JONGER AS 1 MAAND GEBRUIK NIE.
- Onttrekkingsperiodes:
 - Melk: 2 dae
 - Vleis: 7 dae
- Tekens van kwyling en lomerigheid mag voorkom met oordosering. Hierdie tekens is tydelik en diere sal herstel sonder enige nuwe-effekte.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.
- **L.W.** Piretroïedopgiemiddels mag irritasie veroorsaak. Behandel slegs 'n klein groep en neem waar vir 'n uur voordat die res van die kudde behandel word.

VOORSORGMATREËLS

- Moenie eet, drink of rook tydens of na aanwending van die middel alvorens die hande en gesig deeglik gewas is nie.
- Vermoedlik met opgiemiddel en was onmiddellik en deeglik met seep en warm water na toevallige besmetting.
- Verwyder en was besoedelde klere.
- Vernietig leë houers en moenie vir enige ander doel gebruik nie.
- Moenie damme, riviere en ander waterbronne besoedel nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- **Dose:** 1 ml / 10 kg. Treat strictly according to body mass.
- ERADITICK® CATTLE POUR-ON is a ready-to-use product and must not be mixed with any other dip, solvent or insecticidal compound.
- Apply in two lines along each side of the spine from the shoulder to the root of the tail.
- **In cases of heavy brown ear or blue tick challenge,** apply ERADITICK® CATTLE POUR-ON in two lines from the base of the skull, along each side of the spine to the root of the tail.
- **In cases of heavy red-legged tick infestation,** it may be necessary to spot-treat the affected area under the tail above the anus.
- **Treat at regular intervals according to tick challenge.** The longer the product is regularly applied, the more the tick population is reduced. It will then become possible to apply at intervals greater than 7 days.

How to use the applicator

Remove the cap of the calibrated chamber. Without removing the remaining cap, squeeze the container until the correct dosage volume is present in the calibrated chamber, and apply as prescribed

PRESENTATION

200 ml, 500 ml, 1 L, 5 L + 1 L, 20 L

REGISTRATION HOLDER:

Afrivet Business Management (Pty) Ltd
Co. Reg 2000/011263/07
PO Box 2009, Faerie Glen
0043, Pretoria, RSA
Helpline: 0860 833 2787
www.afrivet.co.za

- SLEGS VIR UITWENDIGE GEBRUIK OP BEESTE
Dosis: 1 ml /10 kg. Behandel streng volgens liggaamsmassa.
- ERADITICK® BEES OPGIETMIDDEL is 'n gereed-vir-gebruik produk en moet met geen ander dip, oplossing of insekdodende samestelling gemeng word nie.
- Dien die nodige dosis toe in twee strepe, vanaf die skouer langs beide kante van die ruggraat tot op die stertwortel.
- **In geval van swaar bruinoor- of bloubosluisias,** wend ERADITICK® BEES OPGIETMIDDEL aan in twee strepe vanaf die kopkroon langs beide kante van die ruggraat tot op die stertwortel.
- **In geval van swaar rooipootbosluis besmetting,** mag kolbehandeling van die geaffekteerde area onder die stert bo die anus nodig wees.
- **Behandel met gereelde tussenposes na gelang van bosluispopulasie.** Hoe langer die produk gereeld gebruik word, hoe groter die daling in die bosluisbevolking. Dit mag dan moontlik word om die tussenpose van 7 dae tussen behandelings te verleng.

Gebruik van toediener

Verwyder die prop van die gekalibreerde maatkamer. Sonder om die ander prop te verwyder, druk die toediener totdat die verlangde dosisvolume in die maatkamer versamel, en dien toe soos voorgeskryf.

AANBIEDING

200 ml, 500 ml, 1 L, 5 L + 1 L, 20 L

REGISTRASIEHOUER:

Afrivet Business Management (Pty) Ltd
Mpy. Reg 2000/011263/07
Posbus 2009, Faerie Glen
0043, Pretoria, RSA
Hulplyn: 0860 833 2787
www.afrivet.co.za