

PACKAGE INSERT: Eraditick® Pig Pour-On

FOR EXTERNAL ANIMAL USE ONLY

ERADITICK® PIG POUR-ON

Reg. No. G4253 Act 36/1947
Namibia S0 V03/18.3.4/724

Ready-to-use emulsifiable concentrate

STORAGE INSTRUCTIONS

Store below 25 °C in dry place

COMPOSITION

Amitraz 2 % m/v

INDICATIONS

Kills sarcoptic mange mites and lice on pigs.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST APPLICATION.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this information available.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advise and notify the registration holder.

PRECAUTIONS

- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash contaminated clothing.
- Do not eat, drink or smoke during or after applying before washing hands and face thoroughly.
- Tightly reseal the container if all the

SLEGS VIR UITWENDIGE DIEREGEBRUIK

ERADITICK® VARK OPGIETMIDDEL

Reg. No. G4253 Wet 36/1947
Namibia S0 V03/18.3.4/724

Gereed-om-te-gebruik emulsifiseerbare konsentraat

BERGINSAANWYSINGS

Bewaar benede 25 °C in n droë plek

SAMESTELLING

Amitras 2 % m/v

INDIKASIES

Dood skurftemyte en luise op varke.

WAARSKUWINGS

- MOENIE DIERE SLAG VIR MENSELIKE VERBRUIK BINNE 7 DAE VANAF DIE LAASTE AANWENDING NIE.
- MOENIE OP PERDE GEBRUIK NIE.
- In geval van vergiftiging raadpleeg 'n dokter en maak hierdie inligting beskikbaar.
- Hou buiten die bereik van kinders, oningelegte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Verwyder kontak met opgietmiddel en was onmiddellik en deeglik met seep en warm water in geval van toevallige besmetting.
- Verwyder en was besoedelde klere.
- Moenie eet, drink of rook gedurende en na toediening alvorens hande en gesig deeglig gewas is nie.
- Maak houer dig toe as die inhoud nie

- contents have not been used.
- Destroy empty container by perforation and flattening and NEVER re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

For external animal use only. ERADITICK® PIG POUR-ON is a ready-to-use product and must not be mixed with any other dip, solvent or insecticidal compound.

Apply to all animals in a group whether visually infested or not.

Animals of all ages, including pregnant and suckling sows, piglets, weaners, growers, finishers and boars may be treated.

Apply the calculated amount of ERADITICK® PIG POUR-ON evenly along the backline BETWEEN THE RUMP AND THE EARS.

In large boars and sows the total amount of ERADITICK® PIG POUR-ON should be spread along the backline and sides to increase the spread of the remedy over the body.

The inside of the ears MUST be treated with a small amount of the calculated dose at each application. This is especially important in sows and boars.

Application rate

The dose will provide 5-6 mg of Amitraz per kg body mass and is calculated as quarter the body mass (kg) in ml e.g.:

Body mass (kg)	Application (ml)
5	1,25
10	2,5
20	5,0
30	7,5
40	10,0
50	12,5
60	15,0
100	25,0

- opgebruik is nie.
- Vernietig leë houers deur te perforeer of plat te druk en moet NOoit vir enige ander doeleindes gebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Slegs vir uitwendige dieregebruik.

ERADITICK® PIG POUR-ON is 'n gereed-om-te-gebruik produk en moet nie met enige ander dip, oplosmiddel of insekdodende middel gemeng word nie. Behandel al die diere in 'n groep al is hulle nie sigbaar besmet nie.

Diere van alle ouerdome, insluitend dragtige- en lakterende sôe, jong varkies, speenvarke, groeier, afronders and bere) kan behandel word.

Die berekende hoeveelheid ERADITICK® PIG POUR-ON moet egalig op die ruglyn TUSSEN DIE KRUIS EN ORE aangewend word. In groot bere en sôe moet die totale hoeveelheid op die ruglyn en langs die sye aangewend word om die verspreiding van die middel oor die lyf te vermeerder.

Die binnekant van die ore MOET tydens aanwending met 'n klein hoeveelheid van die berekende hoeveelheid behandel word. Dit is veral belangrik in sôe en bere.

Aanwending

Die dosis sal 5-6 mg Amitras per kg liggaamsmassa verskaf en is bereken op 'n kwart van die liggaamsmassa (kg) in ml byvoorbeeld:

Liggaamsmassa (kg)	Aanwending (ml)
5	1,25
10	2,5
20	5,0
30	7,5
40	10,0
50	12,5
60	15,0
100	25,0

Initial application

Two applications to the entire herd with a 7-10 day interval.

In the case of severe infestation a third dose may be needed 10 days later.

Follow-up and subsequent applications

Apply to sows twice at weekly intervals within the last month before farrowing.

Apply to boars monthly using two applications at weekly intervals.

Recurrence of infestation in growers to be treated twice at weekly intervals.

All introductions to the unit to be treated three times at weekly intervals whilst in quarantine.

Culling of all badly infested sows and boars must be considered.

As ERADITICK® PIG POUR-ON spreads over the surface of the pig's skin, killing parasites in the more inaccessible areas, there is no need to scrub the pig to remove scales before application.

PRESENTATION

100 mL, 200 mL, 250 mL, 500 mL, 1 ℥, 2 ℥, 5 ℥, 10 ℥, 20 ℥

REGISTRATION HOLDER:

Afrivet Business Management (Pty) Ltd
Co. Reg 2000/011263/07
P O Box 2009, Faerie Glen
0043, Pretoria, R.S.A.
Helpline: 0860 833 2787
www.afrivet.co.za

Aanvanklike aanwending

Twee toedienings met 'n 7-10 dae tussenpose op die hele trop.

In geval van swaar besmetting mag 'n derde dosis 10 dae later nodig wees.

Opvolg en daaropvolgende aanwendings

Behandel sôe twee keer met weeklikse tussenposes gedurende die laaste maand voor jong.

Behandel bere maandeliks met twee aanwendings met weeklikse tussenposes.

Herbesmetting in groeiers moet met twee weekliks aanwendings behandel word.

Alle nuwe toevoegings tot die trop moet drie keer met weeklikse tussenposes gedurende kwarentyn behandel word.

Uitwissing van alle erg besmette sôe en bere moet oorweeg word.

Aangesien ERADITICK® PIG POUR-ON oor die oppervlakte van die vark se vel versprei en parasiete in die ontoeganklike areas dood, is dit onnodig om skubbe deur middle van skrop te verwijder.

AANBIEDING

100 mL, 200 mL, 250 mL, 500 mL, 1 ℥, 2 ℥, 5 ℥, 10 ℥, 20 ℥

REGISTRASIEHOUER:

Afrivet Business Management (Pty) Ltd
Mpy. Reg 2000/011263/07
Posbus 2009, Faerie Glen
0043, Pretoria, R.S.A.
Hulphyn: 0860 833 2787
www.afrivet.co.za