

FOR EXTERNAL ANIMAL USE ONLY

ERADITICK® ULTRA

Reg. No. G3976 (Act 36/1947)

Emulsifiable Concentrate Cattle Dip and Spray

Controls ticks on cattle

- Blue ticks
- *Hyalomma* spp.
- *Amblyomma* spp.
- *Rhipicephalus evertsi*
- *Rhipicephalus appendiculatus*

Controls houseflies.

POISONOUS

COMPOSITION

Chlorfenvinphos	20,0 % m/v
Cymiazol	16,0 % m/v
Cypermethrin	2,5 % m/v

STORAGE INSTRUCTIONS

- Store concentrate in original container away from food and feed.
- Store in a cool, dry, well ventilated place away from ignition and heat.

WARNINGS

- DO NOT USE MILK FROM TREATED ANIMALS FOR HUMAN CONSUMPTION WITHIN 12 HOURS OF LAST APPLICATION.
- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 10 DAYS OF LAST APPLICATION.
- Calves younger than 3 months old may show signs of irritation, especially when dipped with a fresh dipwash.
- Sporadic cases of irritation may occur in dairy cows which may lead to a drop in milk production.
- Extremely toxic to fish.
- Handle with care. Poisonous by contact, swallowing or inhalation. In the case of poisoning consult a doctor and make this label available.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Avoid contact with concentrate. In case of accidental skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.
- Avoid prolonged contact with dipwash or inhalation of

SLEGS VIR UITWENDIGE DIEREGERBUIK

ERADITICK® ULTRA

Reg. No. G3976 (Wet 36/1947)

Emulsifiseerbare Konsentraat Bees Dip- en Sproei

Beheer bosluise op beeste

- Bloubosluise
- *Hyalomma* spp.
- *Amblyomma* spp.
- *Rhipicephalus evertsi*
- *Rhipicephalus appendiculatus*

Beheer huisvlieë.

GIFTIG

SAMESTELLING

Chlorfenvinfos	20,0 % m/v
Simiasol	16,0 % m/v
Sipermetrien	2,5 % m/v

BERGINGSAAWYSINGS

- Berg in oorspronklike houer weg van kos en voer.
- Berg in 'n koel, droë, goed geventileerde plek weg van ontbranding en hitte.

WAARSKUWINGS

- MOENIE MELK VAN BEHANDELDE DIERE VIR MENS-LIKE VERBRUIK BINNE 12 URE NA LAASTE BEHANDELING GEBRUIK NIE.
- MOENIE DIERE VIR MENS-LIKE VERBRUIK SLAG BINNE 10 DAE NA LAASTE AANWENDING NIE.
- Tekens van irritasie mag in kalwers jonger as 3 maande oud voorkom, veral wanneer hulle in 'n vars dipmengsel gedip word.
- Irritasie in melkkoeie mag sporadies voorkom wat kan lei tot velaagte melk produksie.
- Uiters giftig vir visse.
- Hanteer versigtig. Giftig na kontak, inname en inaseming. In geval van vergiftiging raadpleeg 'n dokter en maak hierdie etiket beskikbaar.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Vermoed kontak met konsentraat. In geval van toevallige vel kontak, verwyder klere en was vel deeglik met seep en water.
- Vermoed kontak met dipmengsel vir lang periodes en

spray mist.

- Do not eat, drink or smoke while mixing or during use of dip.
- Prevent contamination of food, food utensils, grazing, drinking water, dams, rivers, etc.
- Destroy empty containers by perforation and flattening. Do not re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- DO NOT MIX WITH ANY OTHER DIP.
- SHAKE WELL BEFORE USE.
- Premix by adding the measured amount of concentrate slowly to double the volume of water and mix thoroughly by stirring.
- Add the premix to the bulk of water in the diptank or spray race sump. Make sure that in case of plunge tanks, the larger volume is poured in at the deep end.
- Mix contents of tank well before commencing dipping or spraying.

RECOMMENDATIONS

- **Spray races:** Mix only enough spray-wash for one treatment and do not allow wash to stand over to the next spraying.
- **Replenishment:** It is recommended that not more than 100 head be dipped or sprayed at one time without replenishing the dip, especially where fouling of the wash or tick-borne diseases are a problem. In order to maintain a satisfactory level of the active ingredients refer to the "Head Count" replenishment rate in the table below.

Hand spraying: Thoroughly wet the animal to the skin, especially under the tail, the ears and between the legs where ticks cluster.

TREATMENT INTERVALS

- Dip every 2 weeks.
- Spray once a week.

Dip/Spray according to tick challenge, unless advised otherwise by your veterinarian.

DILUTION/VERDUNNING

METHOD/METODE	FRESHFILL/VARSVUL		"Head Count" REPLENISHMENT/AANVULLING	
	ERADITICK® ULTRA	WATER	ERADITICK® ULTRA	Number of head dipped
Plunge dip/ Dompeldip	1 L	600 L	900 ml	100
Spray race/Spuitdip	1 L	600 L	500 ml	100
Hand spraying/Handspuit	10 ml ERADITICK® ULTRA per 6 L water			

inaseeming van sproeinewel.

- Moenie eet, drink of rook tydens vermenging of gedurende dip nie.
- Voorkom besoedeling van kos, eetgerei, voer, drinkwater, damme, riviere, ens.
- Vernietig leë houers deur te perforeer en plat te druk. Moenie vir enige ander doel gebruik nie.

GEBRUIKSAANWYSINGS GEBRUIK SLEGS SOOS AANGEDUI

- MOENIE MET ENIGE ANDER DIP MENG NIE.
- SKUD GOED VOOR GEBRUIK.
- Maak 'n voormengsel aan deur die regte hoeveelheid konsentraat by twee maal soveel water te voeg, en deeglik te meng.
- Voeg die voormengsel by die res van die water in die diptenk of spuitgang se houertenk. Maak seker dat die meeste van die mengsel aan die diep kant van 'n diptenk gevoeg word.
- Meng die inhoud van die tenk goed voordat daar met dip of bespuiting begin word.

AANBEVEELINGS

- **Spuitdip:** Meng slegs genoeg dipmengsel vir een behandeling en moenie laat oorstaan nie.
- **Versterking:** Dit word aanbeveel dat nie meer as 100 kop op 'n slag gedip of gespuit word nie sonder om die dip aan te vul nie, veral waar besoedeling van die mengsel of bosluisoordraagbare siektes 'n probleem is. Om 'n genoegsame vlak van die aktiewe bestandele te behou verwys na die "Head Count" aanvulling koers in die tabel.

Handbespuiting: Verseker dat diere tot op die vel benat is, veral onder die stert, die ore en tussen die bene waar bosluise versamel.

BEHANDELING TUSSENPOSES

- Dip elke 2 weke.
- Spuit elke week.

Dip/ spuit volgens bosluis uitdaging, tensy anders aanbeveel deur u veearts.

PRESENTATION

100 ml, 250 ml, 500 ml, 1 L, 5 L, 20 L

REGISTRATION HOLDER:

Afrivet Business Management (Pty)
Ltd Co. Reg. No.2000/011263/07
P O Box 2009
Faerie Glen, 0043, R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Helpline: 0860 833 2787

AANBIEDING

100 ml, 250 ml, 500 ml, 1 L, 5 L, 20 L

REGISTRASIEHOUER:

Afrivet Business Management (Pty)
Ltd Mpy. Reg. No.2000/011263/07
Posbus 2009
Faerie Glen, 0043, R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787