

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIEREGBRUIK

EXPEL PLUS
SHEEP DIP & SHEEP AND GOAT JETTING FLUID
Reg. No. G4027 Act 36/1647

EXPEL PLUS
SKAAPDIP & SKAAP- EN BOK SPUITSTOF
Reg. No. G4027 Act 36/1647

CAUTION

VERSIGTIG

STORAGE INSTRUCTIONS

Store in a cool, dry, well ventilated place away from sources of ignition or heat.
Keep only in original container.
Keep container tightly closed.

BERGINGSAAWYSINGS

Berg in 'n koel, droë, goed geventileerde plek weg van bronne van ontbranding en hitte.
Berg slegs in oorspronklike houër.
Verseker dat houër dig toegemaak is.

INDICATIONS

- Kills red lice on sheep & goats.
- Kills blowfly maggots in sheep.
- Prevents blowfly strike for up to 16 weeks.
- Kills sheep scab mites.

AANWYSINGS

- Dood rooi luise op skape en bokke.
- Dood brommer maaiers op skape.
- Voorkom brommer aanvalle vir tot 16 weke.
- Dood skaapbrandsiektemyte.

COMPOSITION

Ivermectin 3% m/v
Novaluron 2% m/v

SAMESTELLING

Ivermektien 3% m/v
Novaluron 2% m/v

WARNINGS

- WITHDRAWAL PERIOD FOR MEAT: 70 DAYS
- DO NOT USE MILK FROM TREATED ANIMALS FOR HUMAN CONSUMPTION.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

WAARSKUWINGS

- ONTREKKINGSSPERIODE VIR VLEIS: 70 DAE.
- MOENIE MELK VAN BEHANDELDE DIERE VIR MENSLIKE VERBUIK GEBRUIK NIE.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

PRECAUTIONS

- Avoid contact with eyes and skin.
- Do not eat, drink or smoke while handling the product.
- Wash hands after use.
- In the case of poisoning contact a doctor and make this label available.
- Destroy empty containers and never re-use for any other purpose.
- Do not contaminate water sources. Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Drug containers and any residual contents should be disposed of by burying or incinerating as free ivermectin may adversely affect fish and certain water-borne organisms.

VOORSORGMAATREËLS

- Vermo kontak met oë en vel.
- Moenie eet, drink of rook gedurende hantering van die produk nie.
- Was hande na gebruik.
- In geval van vergiftiging raadpleeg 'n dokter en maak hierdie etiket beskikbaar.
- Vernietig leë houers en moenie vir enige ander doeleindes gebruik nie.
- Moenie waterbronne besoedel nie. Studies het getoon dat wanneer ivermektien in aanraking met die grond kom, bind dit onmiddellik en stewig met die grond en raak na 'n ruk onaktief. Medisyne houers en enige residuele inhoud moet veilig weggegooi word deur dit te begrawe of verbrand, aangesien vry ivermektien negatiewe effekte op visse en sekere water organismes kan hê.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
SHAKE WELL BEFORE USE

PLUNGE DIP

Premix the concentrate by adding a smaller amount of water to the required quantity of product and stir continuously.
Add this cream to the diptank and mix thoroughly.

Mix as follows (1 : 1000) e.g.

Plunge dip	Water
Freshfill	1 ℓ to 1 000 ℓ water
Replenishment	1 ℓ to 1 000 ℓ water

Ensure that the animals are immersed for approximately 1 minute with the heads being submerged at least twice in order to ensure that the inside of the ears are thoroughly wetted.

JETTING

Premix the concentrate by adding a smaller amount of water to the required quantity of product and stir continuously.
Add this cream to the bulk of water and mix thoroughly.

Mix as follows (1 : 1 000) e.g.

Jetting	Water
1 ml	1 ℓ
10 ml	10 ℓ
100 ml	100 ℓ

- Apply as a jetting spray at a rate of 3 ℓ diluted product for an average size sheep/goat. Ensure thorough wetting.
- Apply as often as necessary i.e. for prevention of blowfly strike and treatment of blowfly maggots apply every 15 – 16 weeks. Wool growth of at least 6 weeks is needed to ensure an optimal protection period.
- For lambs < 6 months of age, repeat treatment after 1 month.
- One treatment for lice infestation will suffice providing that there is NO contact with infested or untreated animals.

SHEEP SCAB MITES

Two treatments with an interval of 7 – 10 days as a plunge dip.

N.B. Sheep scab is a controlled disease in terms of the Animal Disease Act (Act 35/1984). In terms of the Act, any occurrence or suspected occurrence of sheep scab must be reported immediately to the local State Veterinarian.

Presentation:

100 ml, 200 ml, 500 ml, 1ℓ, 5ℓ, 10ℓ and 20 ℓ

Registration holder:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Helpline: 0860 833 2787

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI
SKUD GOED VOOR GEBRUIK

DOMPELDIP

Maak 'n voormengsel deur 'n kleiner hoeveelheid water met die benodigde hoeveelheid produk te meng en roer aanhoudend.
Voeg hierdie room by die afgemete volume water en meng deeglik.

Meng soos volg (1 : 1 000) bv.:

Dompeldip	Water
Varsvul	1 ℓ tot 1 000 ℓ water
Aanvulling	1 ℓ tot 1 000 ℓ water

Verseker dat die diere vir ongeveer 1 minuut gedompel is, en dat hulle koppe ten minste 2 keer onderdompel word, om deeglike benatting van die binnekant van die ore te verseker.

BESPUITING

Maak 'n voormengsel deur 'n kleiner hoeveelheid water met die benodigde hoeveelheid produk te meng en roer aanhoudend.
Voeg hierdie room by die afgemete volume water en meng deeglik.

Meng soos volg (1 : 1 000) bv.:

Bespuiting	Water
1 ml	1 ℓ
10 ml	10 ℓ
100 ml	100 ℓ

- Wend aan as 'n straalspuit teen 'n koers van 3 ℓ verdunde produk vir 'n gemiddelde grote skaap/bok. Verseker deeglike benatting.
- Wend aan so gereeld soos nodig d.w.s. vir voorkoming van brommer aanvalle en behandeling van brommer maaiers, wend aan elke 15 – 16 weke. Ten minste 6 weke wolgroei is nodig om 'n optimale periode van beskerming te verseker.
- Lammers < 6 maande oud moet na 'n maand 'n tweede behandeling ontvang.
- Een behandeling vir luis besmetting sal voldoen mits daar GEEN kontak met besmette of onbehandelde diere is nie.

SKAAPBRANDSIEKTEMYTE

Twee handelings met 'n tussenpose van 7 – 10 dae as 'n dompeldip.

L.W. Onder die Dieresiektes Wet (Wet 35/1984) is skaapbrandsiekte 'n aanmeldbare siekte. In terme van die Wet moet enige geval of vermoede van skaapbrandsiekte onmiddellik by die naaste Staatsveeartskantoor aangemeld word.

Aanbieding:

100 ml, 200 ml, 500 ml, 1ℓ, 5ℓ, 10ℓ and 20 ℓ

Registrasiehouer:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787