

## Package Insert: Salmonella PVI

FOR ANIMAL USE ONLY

### **SALMONELLA PVI**

Reg. No. G4405 Act 36/1947

Salmonella is a notifiable animal disease in terms of the Animal Diseases Act (Act 35/1984). In terms of the Act: Any occurrence of Salmonella or suspected cases have to be reported immediately to the local State Veterinarian.

### **STORAGE**

Store between 2°C to 8°C in a refrigerator. Do not freeze. Protect from light.

### **COMPOSITION**

*S. typhimurium/S. enteritidis/S. gallinarum* ≥ 2.5 x 10<sup>8</sup> CFU/0.5 ml prior to inactivation

### **INDICATIONS**

For the active immunisation of chickens against Salmolnellosis in commercial and breeder birds. Recommended as a prelay and midlay vaccination.

### **WARNINGS**

Birds for human consumption should not be slaughtered within 21 days after treatment.

\* Three to four weeks prior to primary vaccination, all birds should be subjected to a Salmonella rapid plate test using Salmonella pullorum coloured antigen. All birds must be negative for two consecutive tests. Remove any seropositive salmonella reactor birds from the flock. Thorough cleaning and disinfection of the shed is necessary before placement of the flock. A vaccination program will be successful only when it is done in conjunction with Salmonella testing and a good bio-security program.

\* Do not mix this vaccine with other vaccines or medications.

\* All persons handling and administering the vaccine should wear protective clothing.

\* Do not smoke, eat, or drink while handling the vaccine.

This product contains mineral oil. In case of accidental self-injection, seek medical advice immediately and make the leaflet available to the physician.

\* As with all vaccines, occasional hypersensitivity reactions may occur

\* Keep out of reach of children, uninformed persons, and animals.

\* Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice, and notify the registration holder.

### **PRECAUTIONS**

Do not vaccinate layers & breeders within 4 - 6 weeks of lay.

\* Do not use in clinically sick or debilitated birds.

\* Always vaccinate only healthy birds and keep the vaccinated birds under good managemental practices.

\* Use the entire content once opened.

\* Do not use chemical disinfectants for sterilization.

\* Destroy any unused vaccine and dispose of all the vaccine containers and disposable equipment after use in accordance with National Environmental Management: Waste Act, 2008 (Act No. 59 of 2008).

### **DIRECTIONS FOR USE**

USE ONLY AS DIRECTED.

Dosage

#### **For subcutaneous injection:**

Bring the vaccine to room temperature and shake the vaccine container before use.

Use a 20 / 21 gauge needle and automatic syringe. Inject 0.5 ml subcutaneously at the

point midway between the bird's head and base of the neck under the loose skin fold, or intramuscularly in breast muscle.

Follow usual aseptic techniques.

Consult your veterinarian regarding a vaccination schedule.

### **Suggested vaccination schedule**

<b>Bird Type</b>	<b>Primary</b>	<b>Booster</b>	<b>Midlay</b>
Commercial layers	16 <sup>th</sup> week	20 <sup>th</sup> week	Consult your veterinarian
Layer Breeders	16 <sup>th</sup> week	19 <sup>th</sup> week	
Broiler Breeder	18 <sup>th</sup> week	21 <sup>st</sup> week	

### **PRESENTATION**

200 ml (400 doses), 500 ml (1000 doses)

### **REGISTRATION HOLDER**

Afrivet Business Management (Pty) Limited  
P O Box 2009, FAERIE GLEN 0043, R.S.A.

Tel: +27 12 817 9060

Helpline: 0860 833 2787

Web: [www.afrivet.co.za](http://www.afrivet.co.za)