



FOR MORE INFORMATION: +27 12 817 9060 - www.afrivet.co.za

Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07,

PO Box 2009, Faerie Glen, 0043, RSA, **Tel:** +27 12 817 9060, **Web:** www.afrivet.co.za, **Helpline:** 0860 833 2787.

Highly effective and economic hard surface disinfectant
for use in livestock facilities and environment.

A.C.T.

LONG ACTING DISINFECTANT


RSA Reg. No.: NRCS/8054/289982/1189

YOUR PARTNER IN BIOSECURITY

Scientifically proven to eliminate up to **99.999%** of bacteria, fungi and viruses.

PROTECTS YOUR ANIMALS FROM THE MOST
ECONOMICALLY IMPORTANT LIVESTOCK DISEASES INCLUDING
FOOT-AND-MOUTH DISEASE AND AFRICAN SWINE FEVER.

- Residual effect of up to 7 days on surfaces
- Non-corrosive and non-toxic to the environment
- Available in 150 ml (ready-to-use), 1 L, 5 L & 20 L concentrate



RSA Reg. No.: NRCS/8054/289982/1189. **Contains:** Didecylidimethylammonium chloride 0.2 - 0.3 %; benzyl-C12-16-alkyldimethyl chloride 0.2 - 0.3 %; polyhexamethylenebiguanide hydrochloride 0.5 - 0.6 %
Registration holder: RBT24/7, Building C, 5 Regency Drive, Route 21, Irene, Pretoria.
Distributor: Afrivet Business Management (Pty) Ltd, Co. Reg. No. 2000/011263/07.
PO Box 2009, Faerie Glen, 0043, RSA, **Tel:** +27 12 817 9060. **Web:** www.afrivet.co.za, **Helpline:** 0860 833 2787.
FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

 **Afrivet**
Animal health is in our DNA
A Bimeda company

LIVESTOCK SALES TEAM

ANTIBIOTICS:

| | |
|--------------------------|------|
| ULTRATET | Pg 2 |
| ULTRATET LA | Pg 2 |
| ULTRATET LA 200 | Pg 2 |
| ECOSULF L.A. | Pg 3 |
| ECO AFTERBIRTH PESSARIES | Pg 3 |

EXTERNAL PARASITE REMEDY:

| | |
|--|------|
| AFRIVET® 5 % REDLINE ACARICIDAL BALLS | Pg 4 |
| AFRIVET® 10 % REDLINE ACARICIDAL BALLS | Pg 4 |
| AFRIVET® REDLINE POUR-ON | Pg 4 |
| DELTAPOR® 5 | Pg 5 |
| DELTAPOR® 10 PLUS | Pg 5 |
| TIXAGON® 2% | Pg 5 |
| ERADITICK® CATTLE POUR-ON | Pg 6 |
| ERADITICK® PIG POUR-ON | Pg 6 |
| ERADITICK® PLUS POUR-ON | Pg 6 |
| ERADITICK® 125 | Pg 7 |
| ERADITICK® 250 | Pg 7 |
| ERADITICK® ULTRA | Pg 7 |
| ERADITICK® GREASE | Pg 8 |
| DELTA FORCE 100 | Pg 8 |
| EXPEL PLUS JETTING FLUID | Pg 8 |
| ECOBASH | Pg 9 |

INTERNAL PARASITE REMEDY:

| | |
|------------------------|-------|
| ECOFLUKE | Pg 9 |
| ECOTEL 2.5 % | Pg 9 |
| ERADIWORM® CONCENTRATE | Pg 10 |
| ERADIWORM® 25 | Pg 10 |
| ERADIWORM® TAPE | Pg 10 |
| ERADIWORM® PLUS | Pg 11 |
| ECOMINTIC 50 | Pg 11 |
| OSTRIDOSE | Pg 11 |
| MOXZIBEN | Pg 12 |
| CLOSECO 7.5 % | Pg 12 |
| ECOMECTIN 1 % | Pg 12 |
| ECOMECTIN PIG PREMIX | Pg 13 |
| ECOMECTIN SHEEP DRENCH | Pg 13 |

VITAMINS / MINERALS:

| | |
|----------------------|-------|
| VIT-AID | Pg 14 |
| OVI-MIN® GEL | Pg 14 |
| OVI-MIN® + CU GEL | Pg 14 |
| BOVI-MIN® GEL | Pg 15 |
| DAIRY-MIN® GEL | Pg 15 |
| DAIRY-MIN® SUPER | Pg 15 |
| ELECTROGUARD® NF GEL | Pg 16 |
| CALCIMAX® RTU | Pg 16 |
| AFRI-VIT P | Pg 17 |
| AFRI-VIT STRESS | Pg 17 |
| FIBREMAX | Pg 17 |

FEED ADDITIVES:

| | |
|-------------------------------|-------|
| CHLORTET FG 200 | Pg 18 |
| TYLECO FG 100 | Pg 18 |
| AIVLOSIN WATER SOLUBLE POWDER | Pg 19 |

Pg 1

VACCINES:

| | |
|--------------------|-------|
| SALMONELLA VACCINE | Pg 20 |
| WIREVAX | Pg 20 |

HEAT STRESS SOLUTION:

| | |
|----------|-------|
| BETAMINT | Pg 21 |
|----------|-------|

FLY CONTROL:

| | |
|--------------------|-------|
| MISTIFLY® GREEN | Pg 22 |
| ERADIFLY® GRANULAR | Pg 22 |

INSECT CONTROL:

| | |
|-------------|-------|
| ICON® 10 CS | Pg 22 |
|-------------|-------|

WOUND TREATMENTS:

| | |
|-------------------|-------|
| EXPEL WOUND OIL | Pg 23 |
| EXPEL WOUND SPRAY | Pg 23 |
| FUTASPRAY | Pg 23 |

SEALANT:

| | |
|---------------------|-------|
| STOCKHOLM TAR SPRAY | Pg 24 |
|---------------------|-------|

DISINFECTANTS:

| | |
|---------------------------------|-------|
| A.C.T. CLEANING FLUID® | Pg 24 |
| A.C.T. LONG ACTING DISINFECTANT | Pg 24 |

EQUIPMENT:

| | |
|-------------|-------|
| THERMOMETER | Pg 25 |
| WEIGHTBAND | Pg 25 |

ADDITIONAL CONTACT INFORMATION

Pg 26

VACCINATION & DRENCHING PROGRAMMES:

| | |
|---|-------|
| VACCINATION PROGRAMME FOR BEEF CATTLE ON VELD | Pg 27 |
| VACCINATION PROGRAMME FOR DAIRY CATTLE | Pg 27 |
| VACCINATION PROGRAMME FOR SHEEP | Pg 28 |
| SMALL STOCK DRENCHING PROGRAMME BASED ON BREEDING CYCLE | Pg 28 |
| VACCINATION PROGRAMME FOR GOATS | Pg 29 |
| VACCINATION PROGRAMME FOR BROILERS | Pg 29 |
| VACCINATION PROGRAMME FOR PIGS | Pg 30 |
| OPTIONAL VACCINES | Pg 30 |

DIP CALIBRATION

Pg 31

AGEING SHEEP & GOATS BY THEIR TEETH

Pg 32

AGEING CATTLE BY THEIR TEETH

Pg 32



LIVESTOCK SALES TEAM

| | | | | | |
|---|--------------|-------------------------------------|-------------------|-------------------|--------------|
| Bruce Collins | 083 327 3478 | Commercial Manager | | | |
| Northern South Africa & Eswatini | | | | | |
| Head Office | 012 817 9060 | Regional Sales Manager and Eswatini | | | |
| Hesmarie Barkhuizen | 082 324 2966 | Fanie Ungerer | 083 404 9464 | Frik van der Walt | 082 432 7900 |
| Hein Sutherland | 083 388 1003 | Thea Pauley | 078 651 4706 | Thato Khunou | 083 641 6849 |
| Central South Africa | | | | | |
| Riaan Vermeulen | 082 896 4420 | Regional Sales Manager and Feedlots | | | |
| Lee-Ann Terblanche | 079 287 2390 | Louis von Bratt | 071 874 4430 | Danie du Toit | 071 018 5910 |
| Jannie de Klerk | 082 564 3139 | Henk Steyn | 082 899 9406 | Derek Coyne | 082 719 8511 |
| Southern South Africa & Namibia | | | | | |
| Ivahn van Niekerk | 082 410 9517 | Regional Sales Manager | | | |
| Benjamin Jansen van Vuuren | 083 666 1364 | Jacques de Villiers | 066 295 9111 | Japie van Rooyen | 079 496 2332 |
| Francois du Toit | 076 868 0492 | Gareth Webster | 072 508 3381 | Jaco Smit | 082 877 0549 |
| Charl Vorster | 084 812 7378 | Christelle Jooste | 0026 481 470 7406 | | |
| PIG & POULTRY | | | | | |
| Head Office | 012 817 9060 | Sales & Marketing Manager | | | |
| EXPORTS | | | | | |
| Ratselane Marumo | 073 595 9774 | Export Manager Southern Africa | | | |

ANIMAL HEALTH IS IN OUR DNA

ULTRATET

ANTIBIOTIC - TETRACYCLINE INJECTABLE

Oxytetracycline hydrochloride 125 mg / ml



Packaging available:
100 ml, 500 ml

 8 ml / 100 kg  4 ml / 50 kg  1 ml / 25 kg  4 ml / 100 kg



RSA Reg. No.: G0296 (Act 36/1947)
Namibia Reg. No.: NSQV03/17.1.2/698

For the treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill and navel-ill in cattle, sheep, goats and pigs and strangles in horses.

Administration: IM, SC, or slow IV.
Withdrawal period - Meat: 7 days.
Milk: 48 hours.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

ULTRATET LA

ANTIBIOTIC - TETRACYCLINE INJECTABLE

Oxytetracycline hydrochloride 230 mg / ml



Packaging available:
100 ml, 500 ml

 1 ml / 10 kg  1 ml / 10 kg  1 ml / 10 kg (over 20 kg)



RSA Reg. No.: G2857 (Act 36/1947)
Namibia Reg. No.: NSQV01/18.1.2/307

For the treatment and control of conditions caused by, or associated with, oxytetracycline-susceptible organisms. Its long action is recommended where for veterinary, practical or economic reasons it is not possible or desirable to handle and treat the animal frequently.

Cattle: For the treatment of tick-borne gallsickness (anaplasmosis), heartwater, footrot, pneumonia, navel-ill and joint-ill.

Sheep and goats: For treatment of heartwater, footrot, pneumonia, navel-ill and joint-ill.

Pigs: For the treatment of footrot, pneumonia, navel-ill and joint-ill.

Administration: IM (SC for pigs <10kg).
Withdrawal period - Meat: 28 days.
Milk: 5 days.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/00835/07

ULTRATET LA 200

ANTIBIOTIC - TETRACYCLINE INJECTABLE

Oxytetracycline dihydrate 200 mg / ml



Packaging available:
100 ml, 500 ml

 1 ml / 10 kg  1 ml / 10 kg  1 ml / 10 kg (over 20 kg)



RSA Reg. No.: G3559 (Act 36/1947)

Cattle: Treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis).

Sheep and goats: For treatment of heartwater, footrot, pneumonia, navel-ill and joint-ill.

Pigs: For the treatment of pneumonia, navel-ill and joint-ill.

Administration: IM, pigs under 20 kg SC.
Withdrawal period - Meat: 28 days.
Milk: 5 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. 2000/011263/07

ECOSULF L.A.

ANTIBIOTIC - SULPHA INJECTABLE

Sodium sulphadimethoxine 40 % m/v



Packaging available:
100 ml, 500 ml



RSA Reg. No.: G3037 (Act 36/1947)
Namibia Reg. No.: NSOV02/17.1.7/705

Cattle, sheep and goats: Footrot, pneumonia, navel-ill and joint- ill.


Horses: Pneumonia, strangles, joint-ill and navel-ill.

Pigs: Pneumonia and joint-ill.

Calves and lambs: Coccidiosis and bacterial scours.

Administration: SC or IV.
Withdrawal period - Meat: 7 days.
Milk: 48 hours.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

Initial dose  1-2 ml / 10 kg  1-2 ml / 10 kg  1-2 ml / 10 kg  1-2 ml / 10 kg

ECO AFTERBIRTH PESSARIES

ANTIBIOTIC - PESSARIES

Oxytetracycline 375 mg; sulphadimidine 125 mg; sulphadiazine 125 mg; sulphathiazole 125 mg; cetrimide 10 mg / pessary



Packaging available:
1 x 6 pessaries



RSA Reg. No.: G3115 (Act 36/1947)
Namibia Reg. No.: NSOV03/17.1.10/700

For the treatment of bacterial infections of the vagina and uterus following difficult birth and/or retained afterbirth in cattle, sheep, goats, pigs and horses.

Withdrawal period - Meat: 7 days.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

 2 pessaries  ½ pessary  1 pessary  2 pessaries



AFRIVET® 5 % REDLINE ACARICIDAL BALLS



EXTERNAL PARASITE REMEDY - ACARICIDAL BALLS

Flumethrin 5 % m/v

RSA Reg. No.: G4228 (Act 36/1947)



For the control of ticks on blesbuck, kudu and antelope. This product is intended to treat animals that have heavy tick infestation or tick damage and which cannot easily be treated by conventional applications e.g. dip, spray or pour-on. It is not recommended as a general, regular treatment.

Packaging available:
100 x balls



Withdrawal period - Meat: 90 days.
Milk: Not for human consumption.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

65 kg - 125 kg = 1 ball; 125 kg - 250 kg = 2 balls

AFRIVET® 10 % REDLINE ACARICIDAL BALLS



EXTERNAL PARASITE REMEDY - ACARICIDAL BALLS

Flumethrin 10 % m/v

RSA Reg. No.: G4229 (Act 36/1947)



For the control of ticks on blesbuck, kudu and antelope. This product is intended to treat animals that have heavy tick infestation or tick damage and which cannot easily be treated by conventional applications e.g. dip, or pour-on. It is not recommended as a general, regular treatment.

Packaging available:
100 x balls



Withdrawal period - Meat: 90 days.
Milk: Not for human consumption.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

125 kg - 250 kg = 1 ball; 250 kg - 500 kg = 2 balls

AFRIVET® REDLINE POUR-ON



EXTERNAL PARASITE REMEDY - POUR-ON

Flumethrin 1 % m/v

RSA Reg. No.: G4245 (Act 36/1947)
Namibia Reg. No.: NSO/VO4/18.3.3/538
Botswana Reg. No.: W1301246



Controls ticks on cattle, sheep, ostriches, and game.
Controls Tsetse fly worry on cattle.
Controls feather mite on ostriches.
Sterilises female ticks.
Easy application.
Residual action.
Non-systemic action.
The product is rainfast.

Packaging available:
200 ml, 500 ml, 1 L, 5 L, 20 L



Withdrawal period - Meat: 7 days.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

1 ml / 10 kg 1 ml / 10 kg 1 ml / 10 kg 1 ml / 10 kg

DELTAPOR® 5

EXTERNAL PARASITE REMEDY - POUR-ON

Deltamethrin 0.5 % m/v; piperonyl butoxide 2.5 % m/v



Packaging available:
200 ml, 500 ml, 1 L, 5 L, 20 L



RSA Reg. No.: G4252 (Act 36/1947)
Namibia Reg. No.: NSOV04/18.3.3/539
Botswana Reg. No.: W1301250

Ready-to-use pour-on for cattle, sheep and goats. Residual action.
Non-systemic action.

Cattle: Controls ticks, stable flies, horn flies, cattle louse flies and nuisance flies e.g. house flies. Kills lice (biting and sucking).
Protects against blackflies.

Sheep and goats: Controls Karoo-paralysis, bont-legged and red-legged ticks.

Withdrawal period - Meat: 7 days.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

 1 ml / 10 kg  1 ml / 5 kg

DELTAPOR® 10 PLUS

EXTERNAL PARASITE REMEDY - POUR-ON

Deltamethrin 1 % m/v; piperonyl butoxide 3 % m/v



Packaging available:
200 ml, 1 L, 5 L, 20 L



RSA Reg. No.: G4255 (Act 36/1947)

Pour-on for the control of ticks as well as nuisance, biting and tsetse flies on cattle.

Withdrawal period - Meat: 14 days.
Milk: 48 hours.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

 1 ml / 10 kg

TIXAGON® 2%

EXTERNAL PARASITE REMEDY - POUR-ON

Amitraz 2 % m/v



Packaging available:
200 ml



RSA Reg. No.: G3575 (Act 36/1947)

Ready-to-use stripe-on for the control of the yellow dog tick.
Only for use on dogs weighing more than 8 kg.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

 2 ml / 10 kg

ERADITICK® CATTLE POUR-ON

EXTERNAL PARASITE REMEDY - POUR-ON

Amitraz 2% m/v



Packaging available:
20 L



RSA Reg. No.: G4254 (Act 36/1947)
Namibia Reg. No.: [NSO]V04/18.3.4/540
Botswana Reg. No.: W1301249

Ready-to-use pour-on for the control of ticks on cattle.

Withdrawal period - Meat: 7 days.
Milk: 2 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 1 ml / 10 kg

ERADITICK® PIG POUR-ON

EXTERNAL PARASITE REMEDY - POUR-ON

Amitraz 2% m/v



Packaging available:
5 L



RSA Reg. No.: G4253 (Act 36/1947)
Namibia Reg. No.: [NSO]V03/18.3.4/724

Ready-to-use emulsifiable concentrate.
Kills sarcoptic mange mites and lice on pigs.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 12,5 ml / 50 kg

ERADITICK® PLUS POUR-ON

EXTERNAL PARASITE REMEDY - POUR-ON

Amitraz 1,5% m/v; deltamethrin 0,5% m/v; piperonyl butoxide 3% m/v



Packaging available:
1 L, 5 L, 20 L



RSA Reg. No.: G4251 (Act 36/1947)
Namibia Reg. No.: [NSO]V04/18.3.9/541

Ready-to-use pour-on for cattle, sheep and goats.
Residual action. Non-systemic action.

Cattle: Easy application along topline. Controls ticks, stable flies, horn flies, cattle louse flies and nuisance flies e.g. house flies. Kills lice (biting and sucking) and mange mites. Protects against blackflies.

Sheep and goats: Controls Karoo-paralysis, bont-legged and red-legged ticks.

Withdrawal period - Meat: 7 days.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 1 ml / 10 kg  1 ml / 5 kg

ERADITICK® 125

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Amitraz 12.5 % m/v



Packaging available:
200 ml, 500 ml, 1 L, 5 L



RSA Reg. No.: G3583 (Act 36/1947)
Namibia Reg. No.: NSO V10/18.3.4/1147
Botswana Reg. No.: W1301247

Cattle: Controls ticks. Kills lice and mange mites.
Sheep and goats: Controls ticks and sheep scab.
Kills itch mites and goat mange mites.

Cattle: Spray: 1 L / 500 L water. Boost with 100 ml / 100 head
after 400 head sprayed.

Plunge (lime stabilized): 1 L / 2500 L water.

Fresh fill (lime replenishment): 1 L / 500 L water.

Sheep and goats: Plunge (fresh fill): 1 L / 375 L water.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

ERADITICK® 250

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Amitraz 25 % m/v



**DOUBLE
CONCENTRATE**

Packaging available:
500 ml, 1 L, 5 L, 10 L



RSA Reg. No.: G4047 (Act 36/1947)
Namibia Reg. No.: NSO V10/18.3.4/1148

Cattle: Controls ticks. Kills lice and mange mites.
Sheep and goats: Controls ticks and sheep scab.
Kills itch mites and goat mange mites.

Cattle: Spray: 1 L / 1000 L water. Boost with 100 ml / 200 head
after 400 head sprayed.

Plunge (total replenishment): 1 L / 5000 L water.

Fresh fill (lime replenishment): 1 L / 1000 L water.

Sheep and goats: Plunge (fresh fill): 1 L / 750 L water.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

ERADITICK® ULTRA

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Chlorfenvinphos 20 % m/v; cymiazole 16 % m/v; cypermethrin 2.5 % m/v



Packaging available:
5 L, 10 L



RSA Reg. No.: G3976 (Act 36/1947)

Emulsifiable concentrate cattle dip and spray.
Controls ticks on cattle: Blue-, bont-, bont-legged-,
red-legged- and brown-ear ticks.
Controls houseflies.

Plunge dip: 1 L / 600 L water.

Spray race: 1 L / 600 L water.

Hand spraying: 10 ml / 6 L water.

Withdrawal period - Meat: 10 days.

Milk: 12 hours.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

ERADITICK® GREASE

EXTERNAL PARASITE REMEDY - GREASE

Deltamethrin 0.1 % m/v; piperonyl butoxide 0.05 % m/v



Packaging available:
500 g, 4 kg



RSA Reg. No.: G3667 (Act 36/1947)
Namibia Reg. No.: NSOV05/18.3.3/406
Botswana Reg. No.: W1301248

Patch treatment for the control of ticks at the site of application on cattle and sheep.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

 Apply directly with brush to tick feeding sites.

DELTAFORCE 100

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Deltamethrin 10 % m/v



Packaging available:
100 ml, 1 L, 5 L



RSA Reg. No.: G4367 (Act 36/1947)

Cattle plunge and spray dip. Controls all tick species, including blue ticks. Controls nuisance and biting flies on cattle, namely, stable flies, house flies, cattle louse flies, black flies, horn flies. Controls screw-worm infestation and tsetse fly.

Spray dip: 1 L / 2000 L water.
Plunge dip: 1 L / 2666 L water.

Withdrawal period - Meat: 21 days.
Milk: 24 hours.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

EXPEL PLUS JETTING FLUID

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Ivermectin 3 % m/v; novaluron 2 % m/v



Packaging available:
5 L



RSA Reg. No.: G4027 (Act 36/1947)

Kills red lice on sheep and goats.
Kills blowfly maggots in sheep.
Kills sheep scab mites.
Prevents blowfly strike for up to 16 weeks.

Sheep scab mites: Two treatments with an interval of 7 - 10 days as a plunge dip.
Sheep and goats: Plunge dip (fresh fill): 1 L / 1000 L water.
Plunge dip (replenishment): 1 L / 1000 L water.
Jetting: 1 ml / 1 L water.

Withdrawal period - Meat: 70 days.
Milk: Not for human consumption.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

ECOBASH

EXTERNAL PARASITE REMEDY - PLUNGE & SPRAY

Cymiazole 17.5 % m/v; cypermethrin 2.5 % m/v



Packaging available:
1 L, 5 L, 20 L



RSA Reg. No.: G3382 (Act 36/1947)
Namibia Reg. No.: NSQV04/18.3.9/535

Emulsifiable concentrate - cattle plunge and spray dip.
Controls ticks, has a detaching effect. Kills lice.
Controls nuisance flies, biting flies and face flies.
Protects against screw-worm infestation in cattle.

Plunge dip (fresh fill): 1.5 L / 1000 L water).
Plunge dip (replenishment): 3 L / 1000 L water).
Spray race: 1.5 L / 1000 L water.
Hand-spraying: 15 ml / 10 L water.

Withdrawal period - Meat: 7 days.
Milk: Zero.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

See above for detailed instructions.

ECOFLUKE

INTERNAL PARASITE REMEDY - DOSE

Triclabendazole 10 % m/v

⊕†



Packaging available:
5 L



RSA Reg. No.: G3383 (Act 36/1947)
Namibia Reg. No.: NSQV04/18.1.1/537

Liver fluke remedy against immature and adult liver fluke and giant liver fluke in cattle, sheep and goats.

Withdrawal period - Meat: 28 days.
Milk: 14 days.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

 12 ml / 100 kg  1 ml / 10 kg

ECOTEL 2.5 %

INTERNAL PARASITE REMEDY - DOSE

Praziquantel 2.5 % m/v

⊕†



Packaging available:
200 ml, 1 L, 5 L



RSA Reg. No.: G3162 (Act 36/1947)
Namibia Reg. No.: NSQV03/18.1/303

Oral, liquid tapeworm remedy against intestinal and hepatic tapeworms in sheep, and tapeworms in cattle and ostriches.

Cattle dosage: Milk tapeworm 2 ml / 10 kg

Sheep dosage: Milk tapeworm: 2 ml / 10 kg
Serrated tapeworm: 2 ml / 10 kg
Narrow tapeworm: 3 ml / 10 kg
Liver tapeworm: 4 ml / 10 kg

Ostriches dosage: Ostrich tapeworm 3 ml / 10 kg

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

 2 ml / 10 kg  2 - 4 ml / 10 kg  3 ml / 10 kg

ERADIWORM[®] CONCENTRATE

INTERNAL PARASITE REMEDY - DOSE

Levamisole hydrochloride 25 % m/v

③†



Packaging available:
500 ml



RSA Reg. No.: G4250 (Act 36/1947)
Namibia Reg. No.: [NSO]V04/148.1.4/298

Roundworm remedy for cattle, sheep, goats and ostriches.
Does not stain wool or mohair.
Dilute before use - 500 ml to 4.5 L of water.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 15 ml / 50 kg  3 ml / 10 kg  12 ml / 10 kg

ERADIWORM[®] 25

INTERNAL PARASITE REMEDY - DOSE

Levamisole hydrochloride 2.5 % m/v

③†



Packaging available:
200 ml, 1 L



RSA Reg. No.: G4249 (Act 36/1947)
Namibia Reg. No.: [NSO]V02/18.1.4/301

Roundworm remedy for cattle, sheep, goats and ostriches.
Does not stain wool or mohair.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 15 ml / 50 kg  3 ml / 10 kg  12 ml / 10 kg

ERADIWORM[®] TAPE

INTERNAL PARASITE REMEDY - DOSE

Levamisole 37.5 mg/ml; praziquantel 18.8 mg/ml

③† ⑧†



Packaging available:
200 ml, 1 L, 5 L, 10 L



RSA Reg. No.: G4244 (Act 36/1947)
Namibia Reg. No.: [NSO]V04/18.1.8/542

Roundworm and tapeworm remedy for cattle,
sheep, goats and ostriches.

Withdrawal period - Meat: 7 days.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 10 ml / 50 kg  2 ml / 10 kg  1 ml / 1 kg

ERADIWORM® PLUS

INTERNAL PARASITE REMEDY - DOSE

Levamisole 3 % m/v; raxofaxanide 3 % m/v

③† ④†



Packaging available:
500 ml, 1 L, 5 L, 10 L



RSA Reg. No.: G4265 (Act 36/1947)

A remedy against roundworms and liver fluke in sheep, goats and cattle and nasal bot in sheep and goats.

Withdrawal period - Meat: 28 days.
Milk: Not for human consumption.

Registration holder:
Burke Animal Health (Pty) Ltd, Co. Reg. No.: 1996/00721/07

 12.5 ml / 50 kg  2.5 ml / 10 kg

ECOMINTIC 50

INTERNAL PARASITE REMEDY - DOSE

Fenbendazole 5 % m/v

②†



Packaging available:
200 ml, 1 L, 5 L, 10 L



RSA Reg. No.: G2200 (Act 36/1947)
Namibia Reg. No.: NSO V02/18.1.1/1146

A remedy against roundworms in cattle, horses and ostriches and roundworms and milk tapeworm in sheep and goats.

***Sheep and goats:**
Roundworms: 1 ml / 10 kg
Milk tapeworms: 2 ml / 10 kg

Withdrawal period - Meat: 7 days.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/00835/07

 1 ml / 10 kg  1 - 2 ml / 10 kg*  3 ml / 10 kg  2 ml / 10 kg

OSTRIDOSE

INTERNAL PARASITE REMEDY - DOSE

Fenbendazole 5 % m/v

②†



Packaging available:
10 L



RSA Reg. No.: G2243 (Act 36/1947)
Namibia Reg. No.: NSO V01/18.1.1/312

Roundworm remedy for ostriches and horses.

Withdrawal period - Meat: 7 days.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/00835/07

 3 ml / 10 kg  2 ml / 10 kg

MOXZIBEN

INTERNAL PARASITE REMEDY - DOSE

Triclabendazole 5 % m/v; praziquantel 3,75 % m/v; moxidectin 0,1 % m/v

Ⓢ† Ⓢ† Ⓢ†



Packaging available:
5 L, 10 L



RSA Reg. No.: G4040 (Act 36/1947)

Ready-to-use deworming drench for sheep for the control of roundworms i.e. wireworm, brown stomach worm, bankrupt worm, long-necked bankrupt worm and large-mouthed, bowel worm as well as liver fluke, giant liver fluke and milk tapeworm. Controls nasal bot larvae infestation.

Persistence: Wireworm and brown stomach worm for 28 days.

Withdrawal period - Meat: 10 days.
Milk: Not for human consumption.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

 1 ml / 5 kg

CLOSECO 7.5 %

INTERNAL PARASITE REMEDY - INJECTABLE

Closantel 7,5 % m/v

Ⓢ†



Packaging available:
500 ml



RSA Reg. No.: G3440 (Act 36/1947)
Namibia Reg. No.: NSQV04/18.1.3/536

Cattle: Liver fluke and roundworm remedy for cattle with residual action against re-infestation of wireworm, hookworm and nodular worm.

Sheep: Controls wireworm for 21 days, nasalworm larvae and hookworm.

Aids in the control of wireworm for 35 days.
Controls adult liver fluke and aids in the control of immature liver fluke.

Administration (Sheep): SC
Administration (Cattle): SC / IM
Withdrawal period - Meat: 30 days.
Milk: Not for human consumption.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.: 1992/00835/07

 1 ml / 30 kg (normal dose) -  1 ml / 15 kg (double dose)  1 ml / 15 kg

ECOMECTIN 1 %

INTERNAL & EXTERNAL PARASITE REMEDY - INJECTABLE

Ivermectin 1 % m/v

Ⓢ†



Packaging available:
20 ml, 50 ml, 500 ml, 1 L



RSA Reg. No.: G2275 (Act 36/1947)
Namibia Reg. No.: NSQV02/18.1.2/1145

For the treatment of roundworm in sheep and cattle, and false bruising and eye worm in cattle.

Kills sucking lice, mange, screw-worm and controls blue ticks and tampans in cattle.

Kills sheep scab mites, controls Australian itch mite and nasal bots (all stages) in sheep.
Kills mange mites on pigs.

Administration: SC.
Withdrawal period - Meat (Cattle and sheep): 21 days.
Meat (Pigs): 28 days.
Milk: Not for human consumption.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.: 1992/00835/07

 1 ml / 50 kg  1 ml / 50 kg  1 ml / 33 kg

ECOMETIN PIG PREMIX



RSA Reg. No.: G3860 (Act 36/1947)

INTERNAL & EXTERNAL PARASITE REMEDY - POWDER

Ivermectin 0.6 % m/v

⊕†



Antiparasitic remedy for pigs. For the treatment and control of gastro-intestinal roundworms, lungworms, lice and mange in adult and growing pigs.

Treat for 7 consecutive days.

< 40 kg: 333 g per 1 ton of finished feed.

40 - 100 kg: 400 g per ton of finished feed.

Adults: 1.67 kg per ton of finished feed.

Withdrawal period - Meat: > 100 kg: 12 days after last treatment.

< 100 kg: 3 days after last treatment.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

Packaging available:
5 kg

See above for detailed instructions.

ECOMETIN SHEEP DRENCH



RSA Reg. No.: G2630 (Act 36/1947)

Namibia Reg. No.: N\$0V99/18.1.2/305

INTERNAL & EXTERNAL PARASITE REMEDY - DOSE

Ivermectin 0.08 % m/v

⊕†



A remedy against roundworms and nasal bot in sheep and goats and itch mite in sheep.

Withdrawal period - Meat: 14 days.

Milk: Not for human consumption.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd, Co. Reg. No.1992/000835/07

Packaging available:
1 L, 5 L

 2,5 ml / 10 kg



VIT-AID

VITAMINS / MINERALS - INJECTABLE

Vitamin A 500 000 IU; vitamin E 50 IU



Packaging available:
100 ml



RSA Reg. No.: G0678 (Act 36/1947)
Namibia Reg. No.: NSQV02/19.1/472

VIT-AID vitamin A injection is a waterbased solution of vitamin A, for the prevention and treatment of vitamin A deficiency in cattle.

Administration: IM.
Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Formulation Technologies CC, Co. Reg. No.: CK90/19937/23.

 1 ml / 250 kg

OVI-MIN® GEL

VITAMINS / MINERALS - ORAL

Vitamin A 50 000 IU; vitamin D, 2 500 IU; vitamin E 500 IU; manganese 200 mg; zinc 600 mg; selenium 20 mg; magnesium 1 000 mg; cobalt 15 mg; lysine 2 000 mg



Packaging available:
200 ml, 1 L, 5 L, 25 L




RSA Reg. No.: V24722 (Act 36/1947)
Namibia Reg. No.: N-FF2897

Vitamin, mineral and amino acid supplement for sheep and goats. **OVI-MIN® GEL** provides essential organic trace minerals, magnesium, amino acid and fat-soluble vitamins to assist in fertility, immunity and growth in sheep and goats.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23

 Lambs: 5 ml / 12 kg; weaning: 10 ml / 25 kg;
adult ewes: 25 ml / 65 kg; adult rams: 45 ml / 100 - 120 kg

OVI-MIN® + CU GEL

VITAMINS / MINERALS - ORAL

Content per 100 ml: Vitamin A 50 000 IU; vitamin D, 2500 IU; vitamin E 500 IU; magnesium 1000 mg; cobalt 15 mg; lysine 2000 mg; and as organic amino acid complexes: manganese 200 mg; zinc 600 mg; selenium 20 mg and copper 100 mg



Packaging available:
1 L, 5 L, 25 L




RSA Reg. No.: V26349 (Act 36/1947)

Vitamin, mineral and amino acid supplement for sheep and goats. **OVI-MIN® + CU GEL** provides essential organic trace minerals, magnesium, amino acid and fat-soluble vitamins to assist in fertility, immunity and growth in sheep and goats.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23

 Lambs: 5 ml / 12 kg; weaning: 10 ml / 25 kg;
adult ewes: 25 ml / 65 kg; adult rams: 45 ml / 100 - 120 kg

BOVI-MIN® GEL

VITAMINS / MINERALS - ORAL

Content per 100 ml: Vitamin A 100 000 IU; vitamin E 100 IU; magnesium 1000 mg; cobalt 5 mg; lysine 2000 mg; and as organic amino acid complexes: manganese 340 mg; zinc 680 mg; selenium 20 mg and copper 180 mg



Packaging available:
1 L, 5 L, 25 L



Calves: 10 - 20 ml / 80 - 150 kg; cows: 60 ml / 450 kg;
bulls: 120 ml / 850 kg



RSA Reg. No.: V24721 (Act 36/1947)
Namibia Reg. No.: N-FF2896

Vitamin, mineral and amino acid supplement for cattle.
BOVI-MIN® GEL provides essential organic trace minerals, magnesium, amino acid and fat-soluble vitamins to support fertility, immunity and growth in beef cattle.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23.

DAIRY-MIN® GEL

VITAMINS / MINERALS - ORAL

Content per 100 ml: Vitamin A 50 000 IU; vitamin E 400 IU; copper 150 mg; manganese 300 mg; zinc 500 mg; magnesium 500 mg; cobalt 5 mg; selenium 20 mg



Packaging available:
5 L, 25 L

Vitamin, mineral and amino acid supplement for dairy cattle.
DAIRY-MIN® GEL provides essential organic trace minerals, magnesium, amino acid and fat-soluble vitamins to assist in fertility, immunity and growth in dairy cattle.



RSA Reg. No.: V24723 (Act 36/1947)
Namibia Reg. No.: N-FF2898

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23



Calves: 15 - 20 ml / 50 - 90 kg; replacement heifers: 55 - 60 ml / 250 - 275 kg;
cows: 100 - 120 ml / 450 - 600 kg; bulls: 165 - 200 ml / 750 - 900 kg

DAIRY-MIN® SUPER

VITAMINS / MINERALS - ORAL

Content per 1 L: Vitamin A 85 000 IU; vitamin E 1200 IU; magnesium 2500 mg; cobalt 25 mg; fulvic acid 780 000 mg; and as organic complexes: manganese 1500 mg; zinc 2500 mg; selenium 100 mg and copper 750 mg



Packaging available:
5 L, 25 L

Vitamin and mineral supplement for dairy cattle.
DAIRY-MIN® SUPER provides essential organic trace minerals from glycinate- and proteinate complexes with magnesium and fat soluble vitamins, enforced with fulvic acid to support the immune system, fertility, growth and milk quality through favourable somatic cell counts.



RSA Reg. No.: V31994 (Act 36/1947)

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23



0.5 ml / kg for 3 successive days

ELECTRO GUARD® NF GEL



RSA Reg. No.: V28622 (Act 36/1947)

VITAMINS / MINERALS - ORAL

Content per 100 ml: Ascorbic acid 500 mg; biotin 3 mg; calcium 600 mg; folic acid 23 mg; magnesium 230 mg; metabolisable energy (dextrose) 250 KJ; moisture 800 mg; niacin 350 mg; pantothenic acid (calcium panthothenate) 105 mg; potassium chloride 950 mg; pyridoxine (vitamin B₆) 90 mg; riboflavin (vitamin B₂) 70 mg; sodium chloride 1300 mg; thiamine (vitamin B₁) 44 mg; vitamin A 110000 IU; vitamin B₁₂ 0.44 mg; vitamin D, 30 000 IU; vitamin E 525 IU; vitamin K, 35 mg



Vitamin, mineral and amino acid supplement for ruminants, game, ostriches, swine and horses. Water soluble gel for the prevention and support of symptoms associated with stress, as a result of feed refusal, rumen and gutstasis, dehydration, diarrhoea, adaptation problems, disease and deviant behaviour in game, ruminants, ostriches, pigs and horses.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:

Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23.

Packaging available:
200 ml, 1L, 5 L, 25 L

Oral treatment: 3 - 4 ml per 10 kg animal, undiluted for 1 to 3 days as required.
Water application: 3 - 4 ml per litre of drinking water for 3 to 5 days as required.

CALCIMAX® RTU



RSA Reg. No.: V32091 (Act 36/1947)

VITAMINS/MINERALS - ORAL

Each litre contains: Calcium 80 g; magnesium 7 g; propylene glycol and monosaccharides 250 g; vitamin D, 20 000 IU



CALCIMAX® RTU is a ready-to-use combination of calcium, magnesium and vitamin D in an energy based combo from propylene glycol and monosaccharides to assist in calcium, magnesium and energy deficits after calving, lambing, farrowing and foaling. Calcium sources are a unique blend from fast-, medium- and slow-release sources.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:

Camelus Grondstowwe CC, Co. Reg. No.: 1996/41164/23

Packaging available:
500 ml

1 ml / kg body mass



AFRI-VIT P



RSA Reg. No.: V28852 (Act 36/1947)
Namibia Reg. No.: N-FF4717

VITAMINS / MINERALS - POWDER

Composition per kg: Vitamin A acetate 40 000 000 IU; vitamin D₃ 10 000 IU; vitamin E (50%) 14 000 IU; vitamin C 45.0 g; vitamin B₁ 5.0 g; vitamin B₂ 6.0 g; vitamin B₃ 3.6 g; vitamin B₆ 1.28 g; vitamin K₁ 2.22 mg; vitamin B₁₂ 20 g; calcium D-pantothenate 2.5 g; dextrose 676 g; glycine 35 g; methionine 26.4 g; arginine 24 g; lysine 24.74 g; threonine 12 g; tryptophane 4.9 g; sodium 16.08 g; chloride 24.56 g; anoxitol dry 10 g; sodium benzoate 5 g

Vitamin and amino acid supplement for poultry, pigs and calves.

AFRI-VIT P contains all the necessary nutrients to help boost disease recovery.



Packaging available:
100 g, 1 kg, 5 kg

0.5 g - 1 g per litre of drinking water for at least 3 - 5 days

Withdrawal period - Meat: Zero.
Milk: Zero.
Eggs: Zero.

Registration holder:
AfriVet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

AFRI-VIT STRESS



RSA Reg. No.: V31551 (Act 36/1947)

VITAMINS / ELECTROLYTE - POWDER

Composition per kg: Betaine 200 000 mg; vitamin C 180 000 mg; sodium chloride 150 000 mg; vitamin A 15 000 000 IU; vitamin E 25 000 mg; vitamin D₃ 2 000 000 IU; pantothenic acid 10 000 mg; potassium chloride 10 000 mg; vitamin B₁ 5 000 mg; vitamin K 4 000 mg; niacin 2 500 mg; vitamin B₆ 2 000 mg; folic acid 500 mg; biotin 40 mg; vitamin B₁₂ 20 mg

Afri-Vit STRESS contains betaine. It is a concentrated water-soluble electrolyte and can be supplemented into feed or into water over periods of excessive heat. Betaine helps to maintain metabolic functions of cells under different types of osmolaric pressure as is experienced under heat stress.



Packaging available:
100 g



Registration holder:
AfriVet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

FIBREMAX



RSA Reg. No.: V26352 (Act 36/1947)

VITAMINS / MINERALS - FEED ADDITIVE

Composition per 100 ml: Polyethylene glycol (PEG) and polyvinylpyrrolidone (PVP) 30 000 mg; propylene glycol and alkalinizing buffering agents 22 000 mg

A water-soluble solution of anti-tannin & terpene inhibitors fortified by energy and buffering support. This limits the digestive and health constraints from tannin and terpenes associated with browsing and consumption of poor value grazing in dry periods and winter.

FIBREMAX supports animal well-being with more efficient use of available plant material.



Packaging available:
1 L, 5 L

200 ml per 1000 L drinking water

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Camelus Grondstowwe CC. Co. Reg. No.: 1996/41164/23

CHLORTET FG 200

FEED ADDITIVE - TETRACYCLINE

Chlortetracycline 200 g / kg (as the stable chlortetracycline calcium complex)



RSA Reg. No.: G3141 (Act 36/1947)



For the treatment of respiratory and other infections caused by organisms susceptible to chlortetracycline in poultry, turkeys, pigs, calves and feedlot cattle. Not for layers and breeding stock.

Poultry / turkey, pigs, cattle and calves: 1,5 - 2 kg per ton of feed for 5 - 7 days.

Withdrawal period - Meat: 7 days.
Milk: Zero.
Eggs: Not for human consumption.

Registration holder:

ECO Animal Health Southern-Africa (Pty) Ltd. Co. Reg. No.: 1992/000835/07

Packaging available:
25 kg

Exact inclusion rate to be determined by a veterinarian.

TYLECO FG 100

FEED ADDITIVES - MACROLIDE ANTIBIOTIC

Tylosin phosphate 10 % m/m



RSA Reg. No.: G2152 (Act 36/1947)



An aid in improving growth rate and/or feed conversion in pigs, beef cattle and broilers and for feed efficiency and egg production in layers.

Withdrawal period - Meat (Pigs): 15 days.
Meat (Broilers): 3 days.

Registration holder:

ECO Animal Health Southern-Africa (Pty) Ltd. Co. Reg. No.: 1992/000835/07

Packaging available:
25 kg

See product insert for detailed instructions.



AIVLOSIN WATER SOLUBLE POWDER

FEED ADDITIVE - MACROLIDE ANTIBIOTIC

Each gram contains 625 mg Tylvalosin (as tylvalosin tartrate) as active substance



RSA Reg. No.: G1598 (Act 36/1947)
Namibia Reg. No.: NSO V02/17.1.4/1144

An aid in the prevention of mycoplasmosis in broiler chickens and replacement pullets.



Prophylaxis: 1 x 40 g sachet to 2 L warm water and mix with 200 L of drinking water.
Treatment: 1 x 40 g sachet to 2 L warm water and mix with 100 L of drinking water.

Withdrawal period - Meat: 5 days.
Eggs: Zero.

Registration holder:
ECO Animal Health Southern-Africa (Pty) Ltd. Co. Reg. No.: 1992/000835/07

Packaging available:
40 g sachet. Each sachet contains 25 g Tylvalosin.

See above for detailed instructions.



SALMONELLA PVI VACCINE



RSA Reg. No.: G4405 (Act 36/1947)

VACCINE - INJECTABLE

Specified Salmonella species with *S. typhimurium* $\geq 2.5 \times 10^8$ CFU; *S. enteritidis* $\geq 2.5 \times 10^8$ CFU; *S. gallinarum* $\geq 2.5 \times 10^8$ CFU; prior to inactivation and an oil adjuvant



For the use in commercial and breeder birds for immunity against salmonellosis in poultry. Recommended as a prelay and midlay vaccination.

Packaging available:
500 ml

Salmonellosis is a controlled animal disease in terms of the Animal Diseases Act (Act 35/1984). In terms of the Act: Any occurrence of Salmonella or suspected cases have to be reported immediately to the local State Veterinarian.

Withdrawal period - Meat: 21 days.

Eggs: Do not vaccinate layers and breeders within 4 - 6 weeks of lay.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

WIREVAX



RSA Reg. No.: G4200 (Act 36/1947)

VACCINE - INJECTABLE

Haemonchus contortus antigen 5 µg / ml



A vaccine to aid in the control of wireworm infestations in sheep.

Packaging available:
250 ml

Administration: SC.
Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

BE PART OF THE

WIREVAX REVOLUTION

Reg. No. G4200 (Act 36/1947)

1ST VACCINE AGAINST WIREWORM IN SOUTH AFRICA

BETAMINT

HEAT STRESS SOLUTION - ORAL

Each litre contains: Betaine 250 g; vitamin c 90 g; potassium chloride 2 g; magnesium chloride hexahydrate 4 g; calcium chloride 40 g; sodium chloride 20 g



RSA Reg. No.: V28610 (Act 36/1947)
Namibia Reg. No.: N-FF4716



Oral solution specifically for rehydration in pigs and poultry suffering from heat stress syndrome due to high environment temperatures and high humidity.

Contains betaine, an osmolyte to aid the cells retain water.

Can be used before, during and after the onset of heat stress.

Prevention: 1 ml to 1 L of water for 3 - 5 days OR 1 L to 1000 L of water for 3 - 5 days.

Control: 2 ml to 1 L of water for 3 - 5 days OR 2 L to 1000 L of water for 3 - 5 days.

Withdrawal period - Meat: Zero.

Milk: Zero.

Registration holder:

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

Packaging available:
1 L

See above for detailed instructions.



MISTIFLY® GREEN



FLY CONTROL - POWDER

VETERINARY COMPLEMENTARY MEDICINE
28.9 combination product: medium chain fatty acids



Ready-to-use dusting powder. Repels stable flies, house flies, horn flies, mosquitoes and midges.

Packaging available:
2 x 3.6 kg bags per tub

This medicine has not been evaluated by the Medicine Control Council. This medicine is not meant to diagnose, treat, cure or prevent any diseases.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See product insert for detailed instructions.

ERADIFLY® GRANULAR

FLY CONTROL - GRANULES

Novaluron 0.2% (Insect growth regulator)



RSA Reg. No.: L10328 (Act 36/1947)

Controls fly nuisance for up to 54 days.
Contains an insect growth regulator.
Breaks the fly life cycle.
Prevents development between larvae and adult fly stages.
Granules for easy application on manure.

Application: Apply at the rate of 26 g / m².
Apply directly to fly breeding sites around round hay bales, dairy barns, cattle feedlots and all other premise areas where flies can breed.

Packaging available:
5 kg, 25 kg

Registration holder:

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

ICON® 10 CS

INSECT CONTROL - SPRAY

Lambda-cyhalothrin 100 g / L



RSA Reg. No.: L7860 (Act36/1947)
Namibia Reg. No.: N-AR 0814

Control of a broad range of crawling and flying insects, e.g. flies, mosquitoes, beetles, cockroaches and ants.
Very low dose required to kill insects with long-lasting (at least up to 4 weeks) residual effect.
Water-based and non-staining.
Micro-encapsulated formulation.

Mosquitos and malaria vector control (indoor residual sprays): 62.5 ml / 10 L water.
Pests: 75 ml / 10 L water.

Packaging available:
62.5 ml sachets, 1 L

Registration holder:
Syngenta South Africa (Pty) Ltd

See above for detailed instructions.

EXPHEL WOUND OIL

WOUND TREATMENT - OIL

Resins and oils 99,39 % m/v; germicide 0,51 % m/v; deltamethrin 0,1 % m/v



Packaging available:
100 ml



RSA Reg. No.: G3245(Act 36/1947)
Namibia Reg. No.: [NSO] V04.18.3.3/482
Botswana Reg. No.: W130581

Screw-worm and wound remedy.
A multi-purpose remedy which effectively controls and prevents screw-worm/blowfly strike, promotes healing and kills ticks present at time of application. Do not use on cats.

Application: Apply in, around and on wounds.
Apply to prevent screw-worm/blowfly strike, reduce bacterial infection and promote healing.
Screw-worm/blowfly strike treatment:
Apply +/- every 4 days until healed.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

EXPHEL WOUND SPRAY

WOUND TREATMENT - SPRAY

Resins and oils 99,39 % m/v; germicide 0,51 % m/v; deltamethrin 0,1 % m/v



Packaging available:
300 ml



RSA Reg. No.: G3245 (Act 36/1947)
Namibia Reg. No.: [NSO] V04.18.3.3/482
Botswana Reg. No.: W130581

Screw-worm and wound remedy.
A multi-purpose remedy which effectively controls and prevents screw-worm/blowfly strike, promotes healing and kills ticks present at time of application.
Do not use on cats.

Application: Apply in, around and on wounds.
Apply to prevent screw-worm/blowfly strike, reduce bacterial infection and promote healing.
Screw-worm/blowfly strike treatment:
Apply +/- every 4 days until healed.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

FUTASPRAY

WOUND TREATMENT - SPRAY

Chlorhexidine gluconate 100 mg; cetrimide 1 g; gentian violet 500 mg



Packaging available:
400 ml



RSA Reg. No.: G2715 (Act 36/1947)

Antibacterial and fungicidal wound spray with gentian violet.
For the treatment of wounds and skin infections.

Application: Clean the wound area of visible dirt.
Apply FUTASPRAY AEROSOL evenly on the whole wound surface.
Repeat daily if necessary.

Withdrawal period - Meat: Zero.
Milk: Zero.

Registration holder:
Clloff Trust, Co. Reg. No.: 1211/85, Malufa Products

See above for detailed instructions.

STOCKHOLM TAR SPRAY



SEALANT - SPRAY

NOT A VETERINARY MEDICINE



Packaging available:
250 ml

Use after dehorning, shearing and docking.
Use on horses hooves.
Repelling moisture after pruning fruit trees.

Directions: Clean the area thoroughly.
Hold the canister upright, spray the application area.

Distributor:
Atrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07

See above for detailed instructions.

A.C.T. CLEANING FLUID®

DISINFECTANT

Tar acids BP 180-250 °C 16 % m/v; tar oils BP 185-300 °C 65 % m/v;
soap (resin + vegetable oils equal 12.5 %) 20 % m/v; carbolic acid (RW) co-efficient 5



Packaging available:
500 ml, 5 L

General inanimate surface disinfectant
for use on farm buildings and equipment.

General disinfection: 1:250 dilution rate.
When infection diseases are present: 1:100 dilution rate.

Registration holder:
Chemical Process Technologies (Pty) Ltd. Co. Reg. No.: 2001/002516/07

See above for detailed instructions.

A.C.T. LONG ACTING DISINFECTANT

DISINFECTANT

Didecyltrimethylammonium chloride 0.2 - 0.3 %; benzyl-C12-16-alkyldimethyl 0.2 - 0.3 %;
polyhexamethylenbiguanide hydrochloride 0.5 - 0.6 %



Packaging available:
150 ml, 1 L, 5 L, 20 L

RSA Reg. No.: NRCS/8054/289982/1189

Highly effective and economic inanimate hard surface disinfectant
for use in livestock facilities and environment.
Biosecurity agent to protect your animals from the
most economically important livestock diseases including
foot-and-mouth disease and African swine fever.

Dilution required:
Foot-and-mouth disease and African swine fever: 1 ml / 15 ml water.
Contact time: 15 minutes.

Registration holder:
Residual Barrier Technology (Pty) Ltd Co. Reg. No.: 2016/499996/07

See above for detailed instructions.

THERMOMETER

EQUIPMENT



A thermometer is an essential part of any farmer's first aid kit. The best time to take an animal's temperature is early in the morning. The normal temperature for cattle, sheep and goats is 38.5°C.

Instructions for use included on packaging.

WEIGHTBAND

EQUIPMENT



Simple, economic, practical and effective way to estimate weight. Knowing the weight of your livestock in order to provide the animal with the correct dose is crucial in the prevention and treatment of disease.

Languages available:
isiZulu and Sesotho.
isiXhosa and Setswana.
English and Portuguese.

Instructions for use included on packaging.



CONTACT DETAILS

COMMERCIAL TEAM

| | | |
|---------------|----------------------|-----------------------------|
| Bruce Collins | Commercial Manager | bruce.collins@afrivet.co.za |
| Anathi Toki | Management Assistant | anathi.toki@afrivet.co.za |

LIVESTOCK

| | | |
|-------------------|-------------------------------------|--------------------------------|
| Head Office | Regional Sales Manager and Eswatini | admin@afrivet.co.za |
| Riaan Vermeulen | Regional Sales Manager and Feedlots | riaan.vermeulen@afrivet.co.za |
| Ivahn van Niekerk | Regional Sales Manager | ivahn.vanniekerk@afrivet.co.za |

PIGS & POULTRY

| | | |
|-------------|---------------------------|---------------------|
| Head Office | Sales & Marketing Manager | admin@afrivet.co.za |
|-------------|---------------------------|---------------------|

COMPANION ANIMALS

| | | |
|------------------|-----------------------|--------------------------------|
| Thalita Weitzman | Business Unit Manager | thalita.weitzman@afrivet.co.za |
|------------------|-----------------------|--------------------------------|

FEEDLOTS

| | | |
|-----------------|-------------------------------------|-------------------------------|
| Riaan Vermeulen | Regional Sales Manager and Feedlots | riaan.vermeulen@afrivet.co.za |
|-----------------|-------------------------------------|-------------------------------|

EXPORTS

| | | |
|------------------|-----------------|--------------------------------|
| Ratselane Marumo | Exports Manager | ratselane.marumo@afrivet.co.za |
|------------------|-----------------|--------------------------------|

AFRIVET TECHNICAL & TRAINING SERVICES

| | | |
|---------------|-----------------|-----------------------------|
| Didi Claassen | General Manager | didi.claassen@afrivet.co.za |
|---------------|-----------------|-----------------------------|

TENDER

| | | |
|--|--|--|
| For quotations contact tenders@afrivet.co.za | | |
|--|--|--|

VACCINATION & DRENCHING PROGRAMMES

VACCINATION PROGRAMME FOR BEEF CATTLE ON VELD

| GROUP: | VACCINE / ORGANISM: | ADMINISTRATION: |
|--------|---------------------------|--|
| Calves | Anthrax | Weaning |
| | Botulism | Weaning, boost 21 days later |
| | Multiclostridial vaccine | 3 months to weaning age, boost 21 days later |
| | Lumpy skin disease | Weaning age |
| | Rift Valley fever (live) | Weaning age |
| | Heartwater | 0 - 21 days |
| | Redwater | 3 - 9 months |
| | Anaplasma | 3 - 9 months |
| | Rabies (ZeroRab) | See package insert |

| | | |
|---------|--|---|
| Heifers | <i>Brucella abortus</i> • S19 ◊ RB51 | • 4 - 7 months of age • 3 months before first breeding season. |
| | Vibrio Respiratory combinations (BVD, IBR, PI3 and BRSV) | 2 months before first breeding |

| | | |
|-------|--------------------------|------------------------------|
| Bulls | Multiclostridial vaccine | Annual booster |
| | Vibrio | 1 - 2 months before breeding |
| | Botulism | Annual revaccination |

| | | |
|------|---------------------------------|--|
| Cows | Multiclostridial vaccine | Annual booster 2 months before calving |
| | Lumpy skin disease | Annual booster in spring |
| | Anthrax | Annual booster (avoid late pregnancy) |
| | Vibrio Respiratory combinations | 1 - 2 months before breeding 1 - 2 months before breeding |
| | <i>Brucella abortus</i> (RB51) | Repeat before breeding (S19 cannot be repeated) |
| | Rabies (ZeroRab) | See package insert |

VACCINATION PROGRAMME FOR DAIRY CATTLE

| GROUP: | VACCINE / ORGANISM: | ADMINISTRATION: |
|--------|---------------------------|-----------------------------------|
| Calves | Heartwater | 0 - 21 days |
| | Rabies (ZeroRab) | Optional at 3 months |
| | Anthrax | 7 months |
| | Botulism | 7 months, boost 21 days later |
| | Multiclostridial | 3 - 7 months, boost 21 days later |
| | Redwater | 3 - 9 months |
| | Anaplasma | 3 - 9 months |
| | Lumpy skin disease | 7 months |
| | Rift Valley fever (live) | 7 months |

| | | |
|---------|---|--|
| Heifers | <i>Brucella abortus</i> • S19 ◊ RB51 | • 4 - 7 months of age ◊ 3 - 8 months before first breeding season |
| | Respiratory combinations (BVD, IBR, PI3 and BRSV) | 7 months and then 1 - 2 months before first breeding season |

| | | |
|-------|------------------|----------------------|
| Bulls | Multiclostridial | Annual booster |
| | Botulism | Annual revaccination |

| | | |
|---------------------------|---|--|
| Cows | Multiclostridial | Booster in dry cows |
| | Botulism | Annual booster |
| | Lumpy skin disease | Annual booster in spring |
| | Anthrax | Annual booster (avoid late pregnancy) |
| | Respiratory combinations (BVD, IBR, PI3 and BRSV) | Annual booster after calving |
| | <i>Brucella abortus</i> (RB51) | After calving before breeding (S19 cannot be repeated) |
| Rabies (ZeroRab) | See package insert | |

FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07.
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

ANIMAL HEALTH IS IN OUR DNA

CATTLEVACC-AD-23-M7-V1

VACCINATION & DRENCHING PROGRAMMES

VACCINATION PROGRAMME FOR SHEEP

| GROUP: | VACCINE / ORGANISM: | ADMINISTRATION: |
|--------|--|--|
| Lambs | Multiclostridial vaccine, including pulpy kidney | Weaning, repeat 21 days later |
| | Anthrax | Weaning |
| | Blue tongue | Weaning (all 3 vaccines) |
| | Rift Valley fever (live) | Weaning |
| | <i>Mannheimia haemolytica</i> (<i>Pasteurella</i>) pneumonia | Weaning or as directed on package insert |
| | Abscesses | Before weaning and give booster as indicated |
| | Orf | From day old |
| | Botulism | Weaning |
| | Heartwater | 0 - 21 days, monitor lambs for reaction |
| | <i>Brucella</i> Rev 1 | Vaccinate ram lambs only (stud) at weaning |
| | WIREVAX <i>Haemonchus contortus</i> | Consult your agent / veterinarian |

| | | |
|---|--|---|
| Ewes | Multiclostridial vaccine, including pulpy kidney and tetanus | Last few weeks of pregnancy |
| | Anthrax | Annual - avoid late pregnancy |
| | Blue tongue | Annual booster in spring but avoid first third of pregnancy |
| | Enzootic abortion <i>Chlamydomphila</i> | Before breeding |
| | Orf | When required |
| | <i>Corynebacterium ovis</i> abscesses | Annual booster before shearing |
| | <i>Mannheimia haemolytica</i> pneumonia | Before danger period e.g. autumn |
| | Botulism | Annual booste |
| WIREVAX <i>Haemonchus contortus</i> | Consult your agent / veterinarian | |

| | | |
|------|---|-----------------------------------|
| Rams | Multiclostridial vaccine, with pulpy kidney | Annual revaccination |
| | Blue tongue | Annual revaccination |
| | <i>Mannheimia haemolytica</i> pneumonia | Before danger period e.g. autumn |
| | WIREVAX <i>Haemonchus contortus</i> | Consult your agent / veterinarian |

SMALL STOCK DRENCHING PROGRAMME BASED ON BREEDING CYCLE

| GROUP: | TIME / PARASITE: | PRODUCT: |
|--------|---|--|
| Rams | 4 - 6 weeks before mating starts Controls roundworms and tapeworms | ERADIWORM[®] TAPE ECOMECTIN 1% |

| | | |
|------|--|--|
| Ewes | 2 - 3 weeks before lambing starts Controls roundworms and tapeworms | ERADIWORM[®] TAPE ECOMECTIN 1% |
| | One month before lambing Controls roundworms and tapeworms | ERADIWORM[®] TAPE |
| | 30 days after lambing Controls roundworms and tapeworms | ERADIWORM[®] TAPE |

| | | |
|-------|--|---|
| Lambs | 30 days after lambing Controls roundworms and tapeworms | ERADIWORM[®] TAPE |
| | 60 days after lambing Controls roundworms and tapeworms | ERADIWORM[®] TAPE ECOTEL 2.5% |
| | At weaning, Controls roundworms and tapeworms | ERADIWORM[®] TAPE |
| | 2 weeks before sending lambs to the market Controls roundworms and nasal worm | ERADIWORM[®] PLUS |

OPTIONAL VACCINES (please consult your veterinarian):

Rabies (**ZeroRab**) According to product insert

Best to determine deworming schedule and monitor product effectivity by faecal reduction egg count tests.

Speak to your Afrivet agent about how WIREVAX can form part of your vaccination programme.



FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07.
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

VACCINATION PROGRAMME

VACCINATION PROGRAMME FOR GOATS

| GROUP: | VACCINE / ORGANISM: | ADMINISTRATION: |
|--------|--|---|
| Kids | Multiclostridial vaccine, including pulpy kidney | Weaning, repeat 21 days later |
| | Anthrax | Weaning |
| | Rift Valley fever (live) | Weaning |
| | <i>Mannheimia haemolytica</i> pneumonia | Weaning or as directed on package insert. |
| | Orf | From day old. |
| | Heartwater | 0 - 21 days, monitor kids for reactions |
| Does | Multiclostridial vaccine, including pulpy kidney and tetanus | Last few weeks of pregnancy |
| | Anthrax | Annual - avoid late pregnancy |
| | Enzootic abortion (<i>Chlamydomphila</i>) | Before breeding |
| | Orf | When required |
| | <i>Mannheimia haemolytica</i> (<i>Pasteurella</i>) pneumonia | Before danger period e.g. winter |
| Rams | Multiclostridial vaccine with pulpy kidney | Annual revaccination |
| | <i>Mannheimia haemolytica</i> pneumonia | Before danger period, e.g. winter |

OPTIONAL VACCINES (please consult your veterinarian):

| | |
|---------------------------|-----------------------------|
| Rabies (ZeroRab) | According to product insert |
|---------------------------|-----------------------------|

VACCINATION PROGRAMME FOR BROILERS

| AGE (DAYS): | DISEASE: | ROUTE: |
|------------------------------|---|------------------------------|
| 0 - 1 Day at hatchery | Newcastle disease, Infectious bronchitis virus, Infectious bursal disease | Coarse spray/ drinking water |
| Day 10 | Newcastle disease, Infectious bronchitis virus | Spray |
| Day 14 | Infectious bursal disease | Drinking water |
| Day 18 | Newcastle disease | Drinking water |

Vaccine programme is based on specific farm challenge.

**ANIMAL HEALTH IS
IN OUR DNA**

FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07,
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

VACCINATION & DRENCHING PROGRAMMES

VACCINATION PROGRAMME FOR PIGS

| GROUP: | AGE: | ADMINISTRATION: |
|-----------------------|--|--|
| PIGLETS | At birth | Colostrum intake very important Drying piglets Tie off umbilical cord Runty piglets: 2 ml ENERGYN orally day 1 and 2 |
| | Day 1 - 3 after birth | Iron injection (intramuscular injection) Ear tattoo (registered ear tattoo: Animal Identification Act) Tail docking (optional) Teeth clipping (optional) Disinfection of instruments Product: i.e. A.C.T LA DISINFECTANT |
| | Day 4 - 6 after birth | Anticoccidial agent |
| | D10 - Weaning | Creep feed (small amounts, frequently) |
| | D10 - D14 after birth D21 after birth | 2 nd iron injection (intramuscular injection) (optional) Vaccine: Porcine circovirus and <i>Mycoplasma hyopneumoniae</i> Deworming |
| DRY SOW UNIT GILTS | 5 weeks and 2 weeks before mating | Vaccinations: parvovirus, leptospirosis and erysipelas |
| | 5 weeks and 2 weeks before farrowing | Vaccinations: <i>Escherichia coli</i> + Clostridium - for piglets |
| | 2 weeks before farrowing | Internal and external parasite control Product: ECOMECTIN 1% Injectable |
| SOWS | 2 weeks before mating (Booster) | Vaccinations: parvovirus, leptospirosis and erysipelas |
| | 2 weeks before farrowing (Booster) | Vaccinations: <i>E.coli</i> + Clostridium - for piglets |
| | 2 weeks before farrowing | Internal and external parasite control Product: ECOMECTIN 1% Injectable |
| BOARS | Every six months | Vaccinations: parvovirus, leptospirosis and erysipelas Internal and external parasite control Product: ECOMECTIN 1% Injectable |

OPTIONAL VACCINES (please consult your veterinarian):

| | |
|----------------|---|
| Sows and boars | Atrophic rhinitis Glässers (<i>Glaesserella parasuis</i>) Porcine circovirus |
| Gilts | Atrophic rhinitis Glässers (<i>Glaesserella parasuis</i>) Porcine circovirus <i>Mycoplasma hyopneumoniae</i> |
| Piglets | Oedema disease Ileitis (<i>Lawsonia intracellularis</i>) |
| Weaners | Glässers (<i>Glaesserella parasuis</i>) <i>Actinobacillus pleuropneumoniae</i> (APP) |
| Growers | Boars: Immunocastration |

FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07.
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

**ANIMAL HEALTH IS
IN OUR DNA**

PIGVACC-AD-23-M7-V1

DIP CALIBRATION

The dip tank must be calibrated before it can be used. The capacity at minimum working level (1,6 m from floor) and the capacity at normal dipping level (20 cm) below jump-off must be marked. The latter is the level at which the dip should be used. There are various methods of calibrating a dip tank:

1. Surface area measurement:

Because the dip tank is not a simple rectangle, an average length and average width must be calculated:

- Measure A = length at full dipping level.
- Measure B = length of floor.

Average length = $A+B / 2$

- Measure C = width of tank at full dipping level.
- Measure D = width of floor.

Average width = $C+D / 2$

- Measure E = depth of tank from normal dipping level.

Now calculate:


Depth of tank (E) x average width x average length = meters x 1000 = capacity of tank in litres at normal dipping level.

2. Flow meter:

Using a flow meter, add water and mark off at 1000 litre intervals.

3. Manual measurement of volume:

Using a 200 litre drum, add water to dip and mark at 1000 litre intervals.



**ANIMAL HEALTH IS
IN OUR DNA**

FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060


Afrivet Business Management (Pty) Ltd. Co. Reg. No.: 2000/011263/07,
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

AGEING SHEEP & GOATS BY THEIR TEETH


Under 1 year (baby teeth)




1 - 1.5 years (2 tooth)




2 years (4 tooth)



2.5 years (6 tooth)




2.5 - 3 years (full mouth)




> 3 years (worn-broken)




Old



AGEING CATTLE BY THEIR TEETH




Under 21 months:
4 pairs of baby, temporary incisors.



Two years:
1 pairs of adult, permanent incisors.



Two to two and a half years:
2 pairs of adult, permanent incisors.
Two and a half years to three years:
3 pairs of adult, permanent incisors.
Three to Three and a quarter years:
4 pairs of adult, permanent incisors.



Six years:
All four pairs of permanent incisors, beginning to wear down.



Seven years:
All four pairs of permanent incisors worn.



Thirteen years:
All four pairs of permanent incisors badly worn and or broken.

FOR MORE INFORMATION: 0860 833 2787 or +27 12 817 9060

Afrivet Business Management (Pty) Ltd, Co. Reg. No.: 2000/011263/07,
PO Box 2009, Faerie Glen, 0043, RSA, Tel: +27 12 817 9060, Web: www.afrivet.co.za.

ANIMAL HEALTH IS IN OUR DNA

AGEETEETH-AD-23-M7-VV1

NOTES

The company provides the information contained herein in good faith but makes no representation as to its comprehensiveness or accuracy. This document is intended only as a guide to the appropriate precautionary handling of the material by a properly trained person using this product. Individuals receiving the information must exercise their independent judgment in determining its appropriateness for a particular purpose.

THE COMPANY MAKES NO REPRESENTATIONS OR WARRANTIES, EITHER EXPRESS OR IMPLIED, INCLUDING WITHOUT LIMITATION ANY WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE WITH RESPECT TO THE INFORMATION SET FORTH HEREIN OR THE PRODUCT TO WHICH THE INFORMATION REFERS. ACCORDINGLY, THE COMPANY WILL NOT BE RESPONSIBLE FOR DAMAGES RESULTING FROM USE OF OR RELIANCE UPON THIS INFORMATION.



Afrivet
Animal health is in our DNA
A Bimeda company

LABORATORY SERVICES

Dip wash sample analysis

The Afrivet laboratory tests the concentration of the active ingredient in dip wash samples taken from dip tanks and spray races. If tick control in a herd is ineffective, the strength of the product in use may be part of the reason. We also offer experienced advice on replenishment rates and the rotation of products.

The laboratory performs faecal egg counts, faecal egg count reduction tests (FECRT), larval culture and coccidian oocyst counts.

Tick identification and acaricide resistance testing

The lab is using the Shaw Larval Immersion Test (SLIT) on larvae hatched from ticks to test resistance to amitraz, chlorfenvinphos and cypermethrin. This assists farmers, reps and veterinarians in identifying resistant strains of ticks, and in the selection of tick control products that are effective.

ANIMAL HEALTH IS IN OUR DNA

FOR MORE INFORMATION: +27 12 817 9060 - www.afrivet.co.za

Afrivet Business Management (Pty) Ltd Co. Reg. No. 2000/011263/07

Posbus 2009, Faerie Glen, 0043 R.S.A. Tel: +27 12 817 9060 Web: www.afrivet.co.za Helpline: +27 860 833 2787.

LAB-AD-23-M7-V1

ANIMAL HEALTH IS IN OUR DNA[®]

